

SUBSTITUTE FORM PTO-1390

U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE

ATTORNEY'S DOCKET NUMBER
00537-190002**TRANSMITTAL LETTER TO THE UNITED STATES
DESIGNATED/ELECTED OFFICE (DO/EO/US)
CONCERNING A FILING UNDER 35 U.S.C. 371**U.S. APPLICATION NO. (If Known, see 37 CFR
1.5) 09/980133INTERNATIONAL APPLICATION NO.
PCT/US00/15396INTERNATIONAL FILING DATE
5 June 2000PRIORITY DATE CLAIMED
4 June 1999TITLE OF INVENTION
NEUROMEDIN B AND SOMATOSTATIN RECEPTOR AGONISTSAPPLICANT(S) FOR DO/EO/US
Barry A. Morgan and Dean Sadat-Aalaei

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. ☒ This is a **FIRST** submission of items concerning a filing under 35 U.S.C. 371.
2. ☐ This is a **SECOND** or **SUBSEQUENT** submission of items concerning a filing under 35 U.S.C. 371.
3. ☐ This is an express request to promptly begin national examination procedures (35 U.S.C. 371(f)).
4. ☐ The US has been elected by the expiration of 19 months from the priority date (PCT Article 31).
5. ☒ A copy of the International Application as filed (35 U.S.C. 371(c)(2))
 - a. ☒ is attached hereto (required only if not communicated by the International Bureau).
 - b. ☐ has been communicated by the International Bureau.
 - c. ☐ is not required, as the application was filed in the United States Receiving Office (RO/US).
6. ☐ An English language translation of the International Application as filed (35 U.S.C. 371(c)(2)).
7. ☐ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3))
 - a. ☐ are attached hereto (required only if not communicated by the International Bureau).
 - b. ☐ have been communicated by the International Bureau.
 - c. ☐ have not been made; however, the time limit for making such amendments has NOT expired.
 - d. ☐ have not been made and will not be made.
8. ☐ An English language translation of amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).
9. ☐ An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)).
10. ☐ An English language translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).

Items 11 to 16 below concern other documents or information included:

11. ☐ An Information Disclosure Statement under 37 CFR 1.97 and 1.98.
12. ☐ An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
13. ☐ A FIRST preliminary amendment.
☐ A SECOND or SUBSEQUENT preliminary amendment.
14. ☐ A substitute specification.
15. ☐ A change of power of attorney and/or address letter.
16. ☒ Other items or information:
 - ☒ International Preliminary Examination Report
 - ☐
 - ☐
 - ☐
 - ☐

CERTIFICATE OF MAILING BY EXPRESS MAIL

Express Mail Label No.

E124547026US

I hereby certify under 37 CFR §1.10 that this correspondence is being deposited with the United States Postal Service as Express Mail Post Office to Addressee with sufficient postage on the date indicated below and is addressed to the U.S. Patent and Trademark Office, P.O. Box 2327, Arlington, VA 22202.

11-27-01
Date of Deposit

Signature

Barry Jenkins
Typed Name of
Person Signing

U.S. APPLICATION NO. (IF KNOWN) 09/980133		INTERNATIONAL APPLICATION NO. PCT/US00/15396		ATTORNEY'S DOCKET NUMBER 00537-190002	
17. <input checked="" type="checkbox"/> The following fees are submitted: Basic National Fee (37 CFR 1.492(a)(1)-(5)): Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO and International Search Report not prepared by the EPO or JPO..... \$1040 International preliminary examination fee (37 CFR 1.482) not paid to USPTO but International Search Report prepared by the EPO or JPO \$890 International preliminary examination fee (37 CFR 1.482) not paid to USPTO but international search fee (37 CFR 1.445(a)(2)) paid to USPTO..... \$740 International preliminary examination fee paid to USPTO (37 CFR 1.482) but all claims did not satisfy provisions of PCT Article 33(1)-(4)..... \$710 International preliminary examination fee paid to USPTO (37 CFR 1.482) and all claims satisfied provisions of PCT Article 33(1)-(4) \$100 <div style="text-align: right;">ENTER APPROPRIATE BASIC FEE AMOUNT =</div>				CALCULATIONS PTO USE ONLY	
Surcharge of \$130 for furnishing the oath or declaration later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(e)).				\$0.00	
Claims	Number Filed	Number Extra	Rate		
Total Claims	26 - 20 =	6	x \$18	\$108.00	
Independent Claims	1 - 3 =		x \$84	\$0.00	
MULTIPLE DEPENDENT CLAIMS(S) (if applicable)			+ \$280	\$0.00	
TOTAL OF ABOVE CALCULATIONS =				\$998.00	
<input type="checkbox"/> Applicant claims small entity status. See 37 CFR 1.27. The fees indicated above are reduced by 1/2.				\$0.00	
SUBTOTAL =				\$998.00	
Processing fee of \$130 for furnishing the English Translation later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(f))				\$0.00	
TOTAL NATIONAL FEE =				\$0.00	
Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31) \$40.00 per property +				\$0.00	
TOTAL FEES ENCLOSED =				\$998.00	
				Amount to be refunded:	\$
				Charged:	\$
a. <input checked="" type="checkbox"/> A check in the amount of \$998.00 to cover the above fees is enclosed. b. <input type="checkbox"/> Please charge my Deposit Account No. 06-1050 in the amount of \$0.00 to cover the above fees. A duplicate copy of this sheet is enclosed. c. <input checked="" type="checkbox"/> The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. 06-1050. A duplicate copy of this sheet is enclosed.					
NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b) must be filed and granted to restore the application to pending status.					
SEND ALL CORRESPONDENCE TO:					
Y. Rocky Tsao FISH & RICHARDSON P.C. 225 Franklin Street Boston, Massachusetts 02110-2804 (617) 542-5070 phone (617) 542-8906 facsimile			<div style="text-align: center;"> SIGNATURE: </div> <div style="text-align: center;"> Y. Rocky Tsao NAME </div> <div style="text-align: center;"> 34,053 REGISTRATION NUMBER </div>		

NEUROMEDIN B AND SOMATOSTATIN RECEPTOR AGONISTSBACKGROUND OF THE INVENTION

5 The mammalian bombesin (Bn)-related peptides, gastrin-releasing peptide (GRP) and neuromedin B (NMB) have a wide range of biological and pharmacological effects. These include stimulation of the release of numerous gastrointestinal hormones and peptides, stimulation of exocrine gland secretion chemotaxis, contraction of smooth muscle, effects in the central nervous system such as thermoregulation, behavioral effects,
10 maintenance of circadian rhythm, inhibition of TSH release and satiety. Bn-related peptides also function as a growth factor in numerous normal cells (e.g., bronchial cells, endometrial stromal cells and 3T3 cells) as well as neoplastic cells such as human small cell lung cancer cells, rat hepatocellular tumor cells, prostatic cells and breast adenocarcinoma cells.

15 Recent structure-function and cloning studies demonstrate that at least two classes of receptors mediate the actions of Bn-related peptides. One class, the GRP-preferring subtype (GRP receptor or GRP-R), has a high affinity for GRP and low affinity for NMB, whereas the other class, the NMB-preferring subtype (NMB receptor or NMB-R), has a high affinity for
20 NMB and lower affinity for GRP. Both classes of receptors are widely present both in the central nervous system and in the gastrointestinal tract. Until recently, the physiological importance of Bn-related peptides in mediating various processes or which receptor subtype mediated the various reported biological effects of Bn-related peptides was unclear.

25 Five different classes of Bn-receptor antagonists have been described. Jensen, R. T. et al. *Trends Pharmacol. Sci.* 12:13 (1991). Members of a number of these classes have high potency, long duration of action and selectivity for the GRP receptor and thus are useful even *in vivo* for defining the role of GRP or GRP receptors in mediating various

- 2 -

physiological events. However, at present few antagonists for the NMB receptor which are sufficiently selective or potent have been described. (See, e.g., Coy, D., and Taylor, J., U.S. Patent 5,462,926.) Further, NMB has been implicated in the inhibition of lung cancer and gliomas, Cancer Res 1991 Oct 1 51:19 5205-11; J Cell Biochem Suppl 1996 24: 237-46, Peptides 1995 16:6 1133-40; J Pharmacol Exp Ther 1992 Oct 263:1 311-7), stimulation of appetite, (Eur J Pharmacol 1994 Dec 12 271:1 R7-9; Am J Physiol 1997 Jan 272:1 Pt 2 R433-7; Pharmacol Biochem Behav 1996 Aug 54:4 705-11), stimulation of TSH secretion, (hypothyroidism), (Regul Pept 1996 Nov 14 67:1 47-53), and inhibition of aldosterone secretion, (hyperaldosteronism), (Histol Histopathol 1996 Oct 11:4 895-7). Thus, the compounds of the present invention are useful in the investigation of the physiological role played by NMB, and in the development of therapeutic compositions for treatment of NMB-related indications.

As is known in the art, agonists and antagonists of somatostatin are useful for treating a variety of medical conditions and diseases, such as inhibition of H. pylori proliferation, acromegaly, restenosis, Crohn's disease, systemic sclerosis, external and internal pancreatic pseudocysts and ascites, VIPoma, nesidoblastosis, hyperinsulinism, gastrinoma, Zollinger-Ellison Syndrome, diarrhea, AIDS related diarrhea, chemotherapy related diarrhea, scleroderma, Irritable Bowel Syndrome, pancreatitis, small bowel obstruction, gastroesophageal reflux, duodenogastric reflux and in treating endocrinological diseases and/or conditions, such as Cushing's Syndrome, gonadotropinoma, hyperparathyroidism, Graves' Disease, diabetic neuropathy, Paget's disease, and polycystic ovary disease; in treating various types of cancer such as thyroid cancer, hepatome, leukemia, meningioma and conditions associated with cancer such as cancer cachexia; in the treatment of such conditions as hypotension such as orthostatic hypotension and postprandial hypotension and panic attacks; GH

- 3 -

secreting adenomas (Acromegaly) and TSH secreting adenomas. Activation of type 2 but not type 5 subtype receptor has been associated with treating prolactin secreting adenomas. Other indications associated with activation of the somatostatin subtypes are inhibition of insulin and/or glucagon and more particularly diabetes mellitus, hyperlipidemia, insulin insensitivity, Syndrome X, angiopathy, proliferative retinopathy, dawn phenomenon and Nephropathy; inhibition of gastric acid secretion and more particularly peptic ulcers, enterocutaneous and pancreaticocutaneous fistula, Dumping syndrome, watery diarrhea syndrome, acute or chronic pancreatitis and gastrointestinal hormone secreting tumors; inhibition of angiogenesis, treatment of inflammatory disorders such as arthritis; chronic allograft rejection; angioplasty; preventing graft vessel and gastrointestinal bleeding. Somatostatin agonists can also be used for decreasing body weight in a patient. Accordingly, the compounds of the instant invention are useful for the foregoing methods.

Recently, it was reported that a native somatostatin (SS), somatostatin-14 (SS-14), inhibited the cross-linking of ^{125}I -GRP to a 120 kD protein in triton extracts of 3T3 cells and human small cell lung cancer cells which are known to possess bombesin receptors. Recent studies have also demonstrated SS-14 could also weakly inhibit binding to opiate receptors, and subsequent structure-function led to the identification of various D-amino acid-substituted and constrained amino acid-substituted cyclo somatostatin analogs that functioned as potent mu opioid receptor antagonists.

All patents and publications mentioned herein are hereby incorporated by reference in their entirety.

SUMMARY OF THE INVENTION

The present invention relates to a series of analogues having unique structural features, and to a method of selectively modulating biochemical

- 4 -

activity of cells induced by somatostatin and/or neuromedin B.

In one aspect the present invention is directed to a compound of the formula (I),



5

(I)

or a pharmaceutically acceptable salt thereof,

wherein

the α -nitrogen of AA^1 , AA^2 , AA^3 , AA^{3b} , AA^4 , AA^5 , AA^6 , AA^7 , AA^{7b} , and AA^8 each is, independently, optionally substituted with (C_{1-4}) alkyl, (C_{3-4}) alkenyl,

10 (C_{3-4}) alkynyl, or (C_{1-6}) alkyl-C(O)-;

AA^1 is absent or the D- or L-isomer of an amino acid selected from the group consisting of R^{11} , Aac, Aic, Arg, Asn, Asp, Dip, Gln, Glu, Hca, Hyp, Lys, Mac, Macab, Orn, Pro, Ser, Ser(Bzl), Thr, Thr(Bzl), Pip, hArg, Bip, Bpa, Tic, Cmp, Inc, Inp, Nip, Ppc, Htic, Thi, Tra, Cmpi, Tpr, Iia, Alla, Aba, Gba, Car, Ipa, Iaa, Inip, Apa, Mim, Thnc, Sala, Aala, Thza, Thia, Bal, Fala, Pala, Dap, Agly, Pgly, Ina, Dipa, Mnf, Inic, C4c, 5-Iqs, Htqa, 4-Mqc, Thn, α -Chpa, Cit, Nua, Pyp and an optionally substituted aromatic α -amino acid;

wherein said optionally substituted aromatic α -amino acid is optionally substituted with one or more substituents each independently selected from the group consisting of halogen, NO_2 , OH, CN, (C_{1-6}) alkyl, (C_{2-6}) alkenyl, (C_{2-6}) alkynyl, (C_{1-6}) alkoxy, Bzl, O-Bzl, and NR^9R^{10} ;

AA^2 is absent or the D- or L-isomer of an amino acid selected from the group consisting of R^{11} , Aic, Arg, Hca, His, Hyp, Pal, F_5 -Phe, Phe, Pro, Trp, and X^0 -Phe Pip, hArg, Bip, Bpa, Tic, Cmp, , Inc, Inp, Nip, Ppc, Htic, Thi, Tra, Cmpi, Tpr, Iia, Alla, Aba, Gba, Car, Ipa, Iaa, Inip, Apa, Mim, Thnc, Sala, Aala, Thza, Thia, Bal, Fala, Pala, Dap, Agly, Pgly, Ina, Dipa, Mnf, Inic, I-Iqc, 3-Iqc, C4c, 5-Iqs, Htqa, 4-Mqc, Thn, α -Chpa, Cit, Nua, and Pyp;

- 5 -

AA³ is the D- or L-isomer of an amino acid selected from the group consisting of Cys, hCys, Pen, Tpa, Tmpa, Mac, Macab, and an optionally substituted aromatic α -amino acid;

wherein said optionally substituted aromatic α -amino acid is optionally

- 5 substituted with one or more substituents selected from the group consisting of halogen, NO₂, OH, CN, (C₁₋₄)alkyl, (C₂₋₄)alkenyl, (C₂₋₄)alkynyl, (C₁₋₄)alkoxy, Bzl, O-Bzl, NR⁹R¹⁰, Pip, hArg, Bip, Bpa, Tic, Cmp, , Inc, Inp, Nip, Ppc, Htic, Thi, Tra, Cmpi, Tpr, , lia, Alla, Aba, Gba, Car, Ipa, laa, Inip, Apa, Mim, Thnc, Sala, Aala, Thza, Thia, Bal, Fala, Pala, Dap, Agly, Pgly, Ina, Dipa, Mnf, Inic,
- 10 I-lqc, 3-lqc, C4c, 5-lqs, Htqa, 4-Mqc, Thn, α -Chpa, Cit, Nua, and Pyp; AA^{3b} is absent or the D- or L-isomer of an amino acid selected from the group consisting of Pal, 4-Pal, His, Arg, Nal, Trp, Bpa, F₅-Phe, Phe, X⁰-Phe, R¹¹, hArg, Bip, Tic, , Htic, Dip, Sala, Aala, Thza, Thia, Bal, Fala, and Pala;

- AA⁴ is a D- or L-isomer of an optionally substituted amino acid or of an
- 15 optionally substituted aromatic α -amino acid;

wherein said optionally substituted amino acid is selected from the group consisting of Trp, Lys, Orn, hLys, *cis*-4-Acha, *trans*-4-Acha, *trans*-4-Amcha, 4-Pip-Gly, N-Met-Trp, β -Met-Trp, His, hHis, hArg, Bip, Tic, Htic, Dip, Sala, Aala, Thza, Thia, Bal, Fala, Pala, and 4-Pip-Ala;

- 20 wherein the side chain amino group of said optionally substituted amino acid is optionally substituted with R³ and R⁴; and

- wherein said optionally substituted aromatic α -amino acid is optionally substituted with one or more substituents each independently selected from the group consisting of halogen, NO₂, OH, CN, (C₁₋₄)alkyl, (C₂₋₄)alkenyl, (C₂₋₄)alkynyl, Bzl, O-Bzl, and NR⁹R¹⁰;
- 25

AA⁵ is absent, R¹¹, Aic, A3c, A4c, A5c, A6c, Abu, Aib, β -Ala, Bpa, Cha, Deg, Gaba, Ile, Leu, Nal, Nie, Pro, Sar, Ser, Ser(Bzl), Thr, Thr(Bzl), Trp, Val, Pal, F₅-Phe, Phe, X⁰-Phe, or an optionally substituted D- or L- isomer of an amino acid selected from the group consisting of 4-Pip-Gly, 4-Pip-

- 6 -

Ala, *cis*-4-Acha, *trans*-4-Acha, *trans*-4-Amcha, hLys, Lys, Orn, hArg, Bip, Tic, , Htic, Dip, Sala, Aala, Thza, Thia, Bal, Fala, and Pala; wherein the side-chain amino group of said optionally substituted amino acid is optionally mono- or di-substituted with R³ and R⁴;

5 AA⁶ is absent or the D- or L-isomer of an amino acid selected from the group consisting of R¹¹, an optionally substituted aromatic α -amino acid, Cys, hCys, Pen, Tpa, Tmpa, Thr, Thr(Bzl), Ser, Ser(Bzl), hArg, Bip, Tic, , Htic, Dip, Sala, Aala, Thza, Thia, Bal, Fala, and Pala;

AA⁷ is absent or the D- or L-isomer of an amino acid selected from the group
10 consisting of R¹¹, an optionally substituted aromatic α -amino acid, A3c, A4c, A5c, A6c, Abu, Aib, Aic, β -Ala, Arg, Cha, Deg, Gaba, Ile, Leu, Nie, Pip, Pro, Sar, Ser, Ser(Bzl), Thr, Thr(Bzl), Val, Tic, Htic, Sala, Aala, Thza, Thia, Bal, Fala, Pala, hArg, Bip, Bpa, Dip, Pal, Sala, and X⁰-Phe;

AA^{7b} is absent or a D- or L-isomer of an amino acid selected from the group
15 consisting of R¹¹, Bpa, Phe, F₅-Phe, X⁰-Phe, Nal, Pro, Ser, Ser(Bzl), Thr, Thr(Bzl), Trp, hArg, Bip, Tic, Htic, Dip, Sala, Aala, Thza, Thia, Bal, Fala, and Pala;

AA⁸ is absent or the D- or L- isomer of an amino acid selected from the group consisting of R¹¹, Maa, Maaab, Thr, Thr(Bzl), Ser, Ser(Bzl), Tyr,
20 Phe(4-O-Bzl), F₅-Phe, and X⁵-Phe, and an optionally substituted aromatic α -amino acid;

R¹ and R² each is, independently, H, E-, E(O)₂S-, E(O)C-, EEOC-, R¹³, or absent;

R³ and R⁴ each is, independently, (C₁₋₁₂)alkyl, (C₂₋₁₂)alkenyl, (C₂₋₁₂)alkynyl,
25 phenyl, naphthyl, phenyl-(C₁₋₆)alkyl, phenyl-(C₂₋₆)alkenyl, phenyl-(C₂₋₆)alkynyl, naphthyl-(C₁₋₆)alkyl, naphthyl-(C₂₋₆)alkenyl, naphthyl-(C₂₋₆)alkynyl, (cyclo(C₃₋₇)alkyl)-(C₁₋₆)alkyl, (cyclo(C₃₋₇)alkyl)-(C₂₋₆)alkenyl, (cyclo(C₃₋₇)alkyl)-(C₂₋₆)alkynyl, heterocyclyl-(C₁₋₄)alkyl, heterocyclyl-(C₂₋₄)alkenyl, heterocyclyl-(C₂₋

- 7 -




alkynyl, 1-adamantyl, 2-adamantyl, 9-fluorenylmethyl, dicyclopropylmethyl, dimethylcyclopropylmethyl, or benzhydryl;

R^5 is $-OR^6$, $-NR^7R^8$, or absent,

wherein each R^6 , R^7 and R^8 is, independently, H, (C_{1-12}) alkyl, (C_{2-12}) alkenyl, (C_{2-12}) alkynyl, phenyl, naphthyl, phenyl- (C_{1-6}) alkyl, phenyl- (C_{2-6}) alkenyl, phenyl- (C_{2-6}) alkynyl, naphthyl- (C_{1-6}) alkyl, naphthyl- (C_{2-6}) alkenyl, naphthyl- (C_{2-6}) alkynyl, 1-adamantyl, 2-adamantyl, 9-fluorenylmethyl, dicyclopropylmethyl, dimethylcyclopropylmethyl, or benzhydryl;


R^9 and R^{10} each is, independently, H, (C_{1-6}) alkyl, (C_{3-4}) alkenyl, (C_{3-4}) alkynyl, 1-adamantyl, or 2-adamantyl;

R^{11} is, independently for each occurrence, a D- or L-amino acid of the formula:




- 8 -

wherein m and n each is, independently, 1, 2, or 3, and p is 0, 1, or 2;
 R^{12} is, independently for each occurrence, an optionally substituted moiety of the formula:



5

R^{13} is a moiety of the formula



wherein q, r, s, and t each is, independently, 0, 1, 2, 3, 4 or 5;
 R^{19} is absent, H, NH_2 , OH, (C_{1-6}) hydroxyalkyl, $N(R^{27}R^{28})$, SO_3H , or an
 10 optionally substituted moiety selected from the group consisting of
 heterocyclyl, phenyl and naphthyl,

wherein the optionally substituted moiety defined for R^{19} is optionally
 substituted with one or more substituents selected, independently for each
 occurrence, from the group consisting of halogen, NO_2 , OH, (C_{1-6}) alkyl,
 15 (C_{2-6}) alkenyl, (C_{2-6}) alkynyl, (C_{1-6}) alkoxy, NH_2 , mono- or di- (C_{1-6}) alkylamino,
 Bzl, and O-Bzl;

R^{20} is O or absent;

R^{21} is (C_{1-6}) alkyl or absent;

- 9 -

R^{22} is N, O, C, or CH;

R^{23} is (C_{1-6}) alkyl or absent;

R^{24} is N, CH, or C;

R^{25} is NH, O, or absent;

5 R^{26} is SO_2 , CO, or CH;

R^{27} and R^{28} each is, independently, H or (C_{1-6}) alkyl;

E is, independently for each occurrence, an optionally substituted moiety selected from the group consisting of (C_{1-12}) alkyl, (C_{2-12}) alkenyl, (C_{2-12}) alkynyl, phenyl, naphthyl, phenyl- (C_{1-6}) alkyl, phenyl- (C_{2-6}) alkenyl, phenyl- (C_{2-6}) alkynyl, 10 naphthyl- (C_{1-6}) alkyl, naphthyl- (C_{2-6}) alkenyl, naphthyl- (C_{2-6}) alkynyl, (cyclo (C_{3-7}) alkyl)- (C_{1-6}) alkyl, (cyclo (C_{3-7}) alkyl)- (C_{2-6}) alkenyl, (cyclo (C_{3-7}) alkyl)- (C_{2-6}) alkynyl, heterocyclyl- (C_{1-4}) alkyl, heterocyclyl- (C_{2-4}) alkenyl, heterocyclyl- (C_{2-4}) alkynyl, 1-adamantyl, 2-adamantyl, dicyclopropylmethyl, dimethylcyclopropylmethyl, 9-fluorenylmethyl, and benzhydryl;

15 wherein the optionally substituted moiety defined for E is optionally substituted with one or more substituents each independently selected from the group consisting of halogen, OH, Bzl, O-Bzl, NO_2 , CN, COOH, and SH;

X^0 is halogen, NO_2 , OH, (C_{1-6}) alkyl, (C_{1-6}) alkoxy, mono- or di- (C_{1-6}) alkylamino, Bzl, O-Bzl, NR^9R^{10} , or CN;

20 X^1 is H, (C_{1-6}) alkyl, (C_{2-6}) alkenyl, (C_{2-6}) alkynyl, indolyl, imidazolyl, 1-naphthyl, 3-pyridyl, optionally ring-substituted benzyl, or a moiety which corresponds to the side-chain group of Arg, Leu, Gln, Lys, Tyr, His, Thr, Trp, Phe, Val, Ala, Lys, or His;

25 wherein said optionally ring-substituted benzyl is optionally substituted with one or more substituents selected from the group consisting of halogen, OH, (C_{1-6}) alkoxy, mono- or di- (C_{1-6}) alkylamino, (C_{1-4}) alkyl, (C_{2-4}) alkenyl, (C_{2-4}) alkynyl, and NR^9R^{10} ;

- 10 -

- X^2 and X^3 each is, independently, H, halogen, OH, =O, =S, (C₁₋₁₂)alkyl, (C₂₋₁₂)alkenyl, (C₂₋₁₂)alkynyl, phenyl, naphthyl, phenyl-(C₁₋₆)alkyl, phenyl-(C₂₋₆)alkenyl, phenyl-(C₂₋₆)alkynyl, naphthyl-(C₁₋₆)alkyl, naphthyl-(C₂₋₆)alkenyl, naphthyl-(C₂₋₆)alkynyl, (cyclo(C₃₋₇)alkyl)-(C₁₋₆)alkyl, (cyclo(C₃₋₇)alkyl)-(C₂₋₆)alkenyl, (cyclo(C₃₋₇)alkyl)-(C₂₋₆)alkynyl, heterocyclyl-(C₁₋₄)alkyl, heterocyclyl-(C₂₋₄)alkenyl, heterocyclyl-(C₂₋₄)alkynyl, 1-adamantyl, 2-adamantyl, dicyclopropylmethyl, or dimethylcyclopropyl methyl;
- X^4 is H, OH, or NH₂; and
- X^5 is halogen, NO₂, CH₃, OH, Bzl or O-Bzl;
- provided that:
- at least six amino acid residues are present;
 - when AA³ is a D- or L-isomer of an amino acid selected from the group consisting of Cys, hCys, Pen, Tpa, or Tmpa, and AA⁶ is a D- or L-isomer of an amino acid selected from the group consisting of Cys, hCys, Pen, Tpa, or Tmpa, then AA³ and AA⁶ are connected by a disulfide bond;
 - when AA¹ or AA³ is a D- or L-isomer of an amino acid selected from the group consisting of Mac or Macab, then AA⁸ is a D- or L-isomer of an amino acid selected from the group consisting of Maa and Maaab, and when AA⁸ is a D- or L-isomer of an amino acid selected from the group consisting of Maa and Maaab, then AA¹ or AA³ is a D- or L-isomer of Mac or of Macab, and AA¹ or AA³ is connected by a disulfide bond with AA⁸;
 - AA² can be D- or L-Hca only when AA¹ is absent;
 - when one of R¹ or R² is E(O)₂S-, E(O)C-, EOOC-, or R¹³, the other is H;
 - when R⁵ is absent, then one of R¹ or R² is also absent, and the N-terminal amino acid and C-terminal amino acid together form an amide bond;
 - when one of X^2 or X^3 is C=O or C=S, the other is absent; and
- said compound of formula (I) is not of the formula:

D-Phe-Tyr-cyclo(D-Cys-D-Trp-Lys-Cys)-Abu-Thr-NH₂;

Ac-Phe-Tyr-cyclo(D-Cys-D-Trp-Lys-Cys)-Abu-Thr-NH₂;

- 11 -

L-4-NO₂-Phe-Tyr-cyclo(D-Cys-D-Trp-Lys-Cys)-Abu-Thr-NH₂;

Ac-L-4-NO₂-Phe-Tyr-cyclo(D-Cys-D-Trp-Lys-Cys)-Abu-Thr-NH₂;

Hca-Tyr-cyclo(D-Cys-D-Trp-Lys-Cys)-Abu-Thr-NH₂;

D-Dip-Tyr-cyclo(Cys-D-Trp-Lys-D-Cys)-Val-Nal-NH₂;

5 D-4-NO₂-Phe-Phe(4-O-Bzl)-cyclo(D-Cys-D-Trp-Lys-Cys)Cha-Nal-NH₂;

or

D-4-NO₂-Phe-cyclo(D-Cys-Phe(4-O-Bzl)-D-Trp-Lys-Cys)-Val-Tyr-NH₂.

In another aspect, this invention is directed to a pharmaceutical composition comprising one or more of a compound of formula (I), as defined hereinabove, and a pharmaceutically acceptable carrier.

10 In yet another aspect, the present invention is directed to a method of eliciting an agonist effect from one or more of a somatostatin and/or neuromedin B subtype receptor in a subject in need thereof, which comprises administering a compound of formula (I), as described

15 hereinabove, to said subject.

In still another aspect, the present invention is directed to a method of eliciting an antagonist effect from one or more of a somatostatin and/or neuromedin B subtype receptor in a subject in need thereof, which comprises administering a compound of formula (I), as described

20 hereinabove, to said subject.

In a further aspect, the present invention is directed to a method of binding one or more somatostatin and/or neuromedin B subtype receptor in a subject in need thereof, which comprises administering a compound of formula (I), as described hereinabove, to said subject.

25 In a still further aspect, the present invention is directed to the use of one or more compounds according to formula I to bind to the neuromedin B receptor or to one or more of the somatostatin receptors, as when performing an in vitro or in vivo assay.

DETAILED DESCRIPTION OF THE INVENTION

- 12 -

One of ordinary skill will recognize that certain substituents listed in this invention may have reduced chemical stability when combined with one another or with heteroatoms in the compounds. Such compounds with reduced chemical stability are not preferred.

5 In general, the compounds of formula (I) can be made by processes which include processes known in the chemical arts for the production of compounds. Certain processes for the manufacture of formula (I) compounds are provided as further features of the invention and are illustrated by the reaction schemes and examples included herein.


10 In the above structural formulas and throughout the instant application, the following terms have the indicated meanings unless expressly stated otherwise:

The term alkyl is intended to include those alkyl groups of the designated length in either a straight or branched configuration. Exemplary
15 of such alkyl groups are methyl, ethyl, propyl, isopropyl, butyl, sec-butyl, tertiary butyl, pentyl, isopentyl, hexyl, isohexyl and the like. When the term C₆-alkyl is included in a definition it is intended to denote a single covalent bond.

The term alkoxy is intended to include those alkoxy groups of the designated length in either a straight or branched configuration. Exemplary
20 of such alkoxy groups are methoxy, ethoxy, propoxy, isopropoxy, butoxy, isobutoxy, tertiary butoxy, pentoxy, isopentoxy, hexoxy, isohexoxy and the like.

The term halogen or halo is intended to include the halogen atoms
25 fluorine, chlorine, bromine and iodine.

The term cycloalkyl is intended to include a mono-cycloalkyl group or a bi-cycloalkyl group of the indicated carbon number known to those of skill in the art.



The term dimethylcyclopropylmethyl refers to the structure


The term aryl is intended to include aromatic rings known in the art, which can be mono-cyclic, bi-cyclic or tri-cyclic, such as phenyl, naphthyl and anthracyl.

- 5 The term heterocycle includes mono-cyclic and bi-cyclic systems having one or more heteroatoms, such as oxygen, nitrogen and/or sulfur. The ring systems may be aromatic, for example pyridine, indole, quinoline, pyrimidine, thiophene (also known as thienyl), furan, benzothiophene, tetrazole, dihydroindole, indazole, N-formylindole, benzimidazole, thiazole, and thiadiazole. The ring systems also may be non-aromatic, for example
- 10 pyrrolidine, piperidine, morpholine and the like.

The chemist of ordinary skill will recognize that certain combinations of heteroatom-containing substituents listed in this invention define compounds which will be less stable under physiological conditions.


- 15 Accordingly, such compounds are less preferred.

- As defined herein, certain residues or moieties are alternatively absent from certain peptides of the invention. Where the bond(s) to such a residue or moiety is indicated by a solid line it is understood that when the residue or moiety is absent a bond is formed between the remaining N-terminal residue or moiety(-ies) and the remaining C-terminal residue or moiety(-ies). Where the bond(s) to such a residue or moiety is indicated by dashed line(s) it is understood that when the residue or moiety is absent no bond is formed between the remaining N-terminal residue or moiety(-ies) and the remaining C-terminal residue or moiety(-ies). For example, in the
- 20 following structure:
- 25




- 14 -


the absence of AA¹ results in




and the absence of AA¹ results in



5 In the following structure:




the absence of R²³ results in



10 When a chemical structure as used herein has an arrow emanating from it, the arrow indicates the point of attachment. For example, the structure



15 is a pentyl group. When a line is drawn through a cyclic moiety, the line indicates that the substituent can be attached to the cyclic moiety at any of the available bonding points. For example,



means that the substituent "X" can be bonded ortho, meta or para to the point of attachment. Similarly, when a line is drawn through a bi-cyclic or a tri-cyclic moiety, the line indicates that the substituent can be attached to the


- 15 -

bi-cyclic or a tri-cyclic moiety at any of the available bonding points in any of the rings.

For all formulas depicted herein the N-terminus is at the left and the C-terminus at the right in accordance with the conventional representation of a polypeptide chain.

The symbol AA¹, AA², or the like in a peptide sequence stands for an amino acid residue, i.e., =N-CH(R)-CO- when it is at the N-terminus or -NH-CH(R)-CO- when it is not at the N-terminus, where R denotes the side-chain of that amino acid residue. Thus, R is -CH(CH₃)₂ for Val. Also, when the amino acid residue is optically active, it is the L-form configuration that is intended unless D-form is expressly designated.

Unless otherwise indicated, where an acetyl group appears at the N-terminus it is understood that the acetyl group is attached to the α -nitrogen rather than to the side chain of the N-terminal amino acid. For example, the structure of the amino acid sequence Ac-4-NO₂-Phe-AA²-AA³- ... is:



Where the substituent Y appears as, e.g., -OR⁵, at the C-terminus of the peptide, it is to be understood that -OR⁵ is attached directly to the carbonyl carbon in replacement of the -OH group.

E(O)C- stands for $\text{---}\overset{\text{O}}{\parallel}\text{C---E}$ and EEOC- stands for $\text{---}\overset{\text{O}}{\parallel}\text{C---O---E}$.


What is meant by "aromatic α -amino acid" is an amino acid residue of

the formula $\text{---NH---}\overset{\text{H}_2\text{C---Z}}{\underset{|}{\text{CH}}}\text{---CO---}$, where Z is a moiety containing an aromatic

- 16 -


ring. Examples of Z include, but are not limited to, a benzene or pyridine ring and the following structures with or without one or more substituent X on the aromatic ring (where X is, independently for each occurrence, halogen, NO₂, CH₃, OH, Bzl, or O-Bzl):

5



Other examples of an aromatic α -amino acid of the invention are substituted His, such as MeHis, His (τ -Me), or His (π -Me).


10 What is meant by nucleic acid base is an optionally substituted nucleic acid moiety of the formula:



15 In certain embodiments of the invention the side chain amino group of one or more amino acids is optionally mono- or di-substituted with R³ and R⁴.

- 17 -

For example, substituting R^3 onto the side chain amino group of 4-Pip-Gly



would result in the following structure: $-NH-CH-CO-$

The compounds of the instant invention have at least one asymmetric center. Additional asymmetric centers may be present on the molecule depending upon the nature of the various substituents on the molecule. Each such asymmetric center will produce two optical isomers and it is intended that all such optical isomers, as separated, pure or partially purified optical isomers, racemic mixtures or diastereomeric mixtures thereof, be included within the scope of the instant invention.

The instant compounds can be generally isolated in the form of their pharmaceutically acceptable acid addition salts, such as the salts derived from using inorganic and organic acids. Examples of such acids are hydrochloric, nitric, sulfuric, phosphoric, formic, acetic, trifluoroacetic, propionic, maleic, succinic, D-tartaric, L-tartaric, malonic, methane sulfonic and the like. In addition, certain compounds containing an acidic function such as a carboxy can be isolated in the form of their inorganic salt in which the counter-ion can be selected from sodium, potassium, lithium, calcium, magnesium and the like, as well as from organic bases.

The pharmaceutically acceptable salts are formed by taking about 1 equivalent of a compound of formula (I) and contacting it with about 1 equivalent of the appropriate corresponding acid of the salt which is desired. Work-up and isolation of the resulting salt is well-known to those of ordinary skill in the art.

The compounds of this invention can be administered by oral, parenteral (e.g., intramuscular, intraperitoneal, intravenous or subcutaneous

- 18 -

injection, or implant), nasal, vaginal, rectal, sublingual or topical routes of administration and can be formulated with pharmaceutically acceptable carriers to provide dosage forms appropriate for each route of administration.

Accordingly, the present invention includes within its scope pharmaceutical compositions comprising, as an active ingredient, at least one of the compounds of formula (I) in association with a pharmaceutically acceptable carrier.

Solid dosage forms for oral administration include capsules, tablets, pills, powders and granules. In such solid dosage forms, the active compound is admixed with at least one inert pharmaceutically acceptable carrier such as sucrose, lactose, or starch. Such dosage forms can also comprise, as is normal practice, additional substances other than such inert diluents, e.g., lubricating agents such as magnesium stearate. In the case of capsules, tablets and pills, the dosage forms may also comprise buffering agents. Tablets and pills can additionally be prepared with enteric coatings.

Liquid dosage forms for oral administration include pharmaceutically acceptable emulsions, solutions, suspensions, syrups, the elixirs containing inert diluents commonly used in the art, such as water. Besides such inert diluents, compositions can also include adjuvants, such as wetting agents, emulsifying and suspending agents, and sweetening, flavoring and perfuming agents.

Preparations according to this invention for parenteral administration include sterile aqueous or non-aqueous solutions, suspensions, or emulsions. Examples of non-aqueous solvents or vehicles are propylene glycol, polyethylene glycol, vegetable oils, such as olive oil and corn oil, gelatin, and injectable organic esters such as ethyl oleate. Such dosage forms may also contain adjuvants such as preserving, wetting, emulsifying, and dispersing agents. They may be sterilized by, for example, filtration through a bacteria-retaining filter, by incorporating sterilizing agents into the

- 19 -

compositions, by irradiating the compositions, or by heating the compositions. They can also be manufactured in the form of sterile solid compositions which can be dissolved in sterile water, or some other sterile injectable medium immediately before use.

- 5 Compositions for rectal or vaginal administration are preferably suppositories which may contain, in addition to the active substance, excipients such as coca butter or a suppository wax.

Compositions for nasal or sublingual administration are also prepared with standard excipients well known in the art.

- 10 Further, a compound of this invention can be administered in a sustained release composition such as those described in the following patents. U.S. Patent No. 5,672,659 teaches sustained release compositions comprising a bioactive agent and a polyester. U.S. Patent No. 5,595,760 teaches sustained release compositions comprising a bioactive agent in a gelable form. U.S. Application No. 08/929,363 filed September 9, 1997, teaches polymeric sustained release compositions comprising a bioactive agent and chitosan. U.S. Application No. 08/740,778 filed November 1, 1996, teaches sustained release compositions comprising a bioactive agent and cyclodextrin. U.S. Application No. 09/015,394 filed January 29, 1998, teaches absorbable sustained release compositions of a bioactive agent. The teachings of the foregoing patents and applications are incorporated herein by reference.

- 25 In general, an effective dosage of active ingredient in the compositions of this invention may be varied; however, it is necessary that the amount of the active ingredient be such that a suitable dosage form is obtained. The selected dosage depends upon the desired therapeutic effect, on the route of administration, and on the duration of the treatment, all of which are within the realm of knowledge of one of ordinary skill in the art.

- 20 -

Generally, dosage levels of between 0.0001 to 100 mg/kg of body weight daily are administered to humans and other animals, e.g., mammals.

A preferred dosage range is 0.01 to 10.0 mg/kg of body weight daily, which can be administered as a single dose or divided into multiple doses, or
5 provided for continuous administration.

Compounds of the instant invention can be and were assessed for their ability to bind to a somatostatin subtype receptor according to the following assays.

The affinity of a compound for human somatostatin subtype receptors 1
10 to 5 (ss_{t1} , ss_{t2} , ss_{t3} , ss_{t4} and ss_{t5} , respectively) is determined by measuring the inhibition of [125 I-Tyr 11]SRIF-14 binding to CHO-K1 cells transfected with the ss_{t1} receptor subtype.

The human ss_{t1} receptor gene was cloned as a genomic fragment. A
1.5 Kb *Pst*I-*Xmn*I segment containing 100 bp of the 5'-untranslated region,
15 1.17 Kb of the entire coding region, and 230 bp of the 3'-untranslated region was modified by the BglII linker addition. The resulting DNA fragment was subcloned into the *Bam*HI site of a pCMV-81 to produce the mammalian expression plasmid (provided by Dr. Graeme Bell, University of Chicago, Chicago, IL.). A clonal cell line stably expressing the ss_{t1} receptor was
20 obtained by transfection into CHO-K1 cells (American Type Culture Collection, Manassas, VA) ("ATCC") using the calcium phosphate co-precipitation method. The plasmid pRSV-neo (ATCC) was included as a selectable marker. Clonal cell lines were selected in RPMI 1640 media (Sigma Chemical Co., St. Louis, MO) containing 0.5 mg/ml of geneticin
25 (Gibco BRL, Grand Island, NY) ring cloned, and expanded into culture.

The human ss_{t2} somatostatin receptor gene, isolated as a 1.7Kb *Bam*HI-*Hind*III genomic DNA fragment and subcloned into the plasmid vector pGEM3Z (Promega), was kindly provided by Dr. G. Bell (University of Chicago, Chicago, IL.). The mammalian cell expression vector is constructed

- 21 -

by inserting the 1.7Kb *Bam*H1-*Hind*III fragment into compatible restriction endonuclease sites in the plasmid pCMV5. A clonal cell line is obtained by transfection into CHO-K1 cells using the calcium phosphate co-precipitation method. The plasmid pRSV-neo is included as a selectable marker.

5 The human *sst*₃ was isolated at genomic fragment, and the complete coding sequence was contained within a 2.4 Kb *Bam*HI/*Hind*III fragment. The mammalian expression plasmid, pCMV-h3 was constructed by inserting the a 2.0 Kb *Nco*I-*Hind*III fragment into the *Eco*R1 site of the pCMV vector after modification of the ends and addition of *Eco*R1 linkers. A clonal cell
10 line stably expressing the *sst*₃ receptor was obtained by transfection into CHO-K1 cells (ATCC) using the calcium phosphate co-precipitation method. The plasmid pRSV-neo (ATCC) was included as a selectable marker. Clonal cell lines were selected in RPMI 1640 media containing 0.5 mg/ml of G418 (Gibco), ring cloned, and expanded into culture.

15 The human *sst*₄ receptor expression plasmid, pCMV-HX was provided by Dr. Graeme Bell (University of Chicago, Chicago, IL.). The vector contains the 1.4 Kb *Nhe*I-*Nhe*I genomic fragment encoding the human *sst*₄, 456 bp of the 5'-untranslated region and 200 bp of the 3'-untranslated region, cloned into the *Xba*I/*Eco*R1 sites of PCMV-HX. A clonal cell line
20 stably expressing the *sst*₄ receptor was obtained by transfection into CHO-K1 cells (ATCC) using the calcium phosphate co-precipitation method. The plasmid pRSV-neo (ATCC) was included as a selectable marker. Clonal cell lines were selected in RPMI 1640 media containing 0.5 mg/ml of G418 (Gibco), ring cloned, and expanded into culture.

25 The human *sst*₅ gene was obtained by PCR using a λ genomic clone as a template, and kindly provided by Dr. Graeme Bell (University of Chicago, Chicago, IL). The resulting 1.2 Kb PCR fragment contained 21 base pairs of the 5'-untranslated region, the full coding region, and 55 bp of the 3'-untranslated region. The clone was inserted into *Eco*R1 site of the

- 22 -

plasmid pBSSK(+). The insert was recovered as a 1.2 Kb *HindIII*-*XbaI* fragment for subcloning into pCVM5 mammalian expression vector. A clonal cell line stably expressing the SST₅ receptor was obtained by transfection into CHO-K1 cells (ATCC) using the calcium phosphate co-precipitation method. The plasmid pRSV-neo (ATCC) was included as a selectable marker. Clonal cell lines were selected in RPMI 1640 media containing 0.5 mg/ml of G418 (Gibco), ring cloned, and expanded into culture.

CHO-K1 cells stably expressing one of the human sst receptors are grown in RPMI 1640 containing 10% fetal calf serum and 0.4 mg/ml geneticin. Cells are collected with 0.5 mM EDTA, and centrifuged at 500g for about 5 minutes at about 4°C. The pellet is resuspended in 50 mM Tris[hydroxymethyl]aminomethane hydrochloride, pH = 7.4 at 25°C, ("Tris buffer"), and centrifuged twice at 500g for about 5 minutes at about 4°C. The cells are lysed by sonication and centrifuged at 39,000g for about 10 minutes at about 4°C. The pellet is resuspended in the same buffer and centrifuged at 50,000g for about 10 minutes at about 4°C and membranes in resulting pellet are stored at - 80°C.

Competitive inhibition experiments of [¹²⁵I-Tyr¹¹]SRIF-14 binding are run in duplicate in polypropylene 96 well plates. Cell membranes (10 µg protein/well) are incubated with [¹²⁵I-Tyr¹¹]SRIF-14 (Dr. Tom Davis, Univ. of Arizona, Tuscon, AZ) (0.05 nM) for about 60 minutes at about 37°C in 50 mM HEPES, 0.2% BSA, 2.5 mM MgCl₂.

Bound from free [¹²⁵I-Tyr¹¹]SRIF-14 is separated by immediate filtration through GF/C glass fiber filter plate (Unifilter, Packard, Meriden, CT) presoaked with 0.3% polyethylenimine (P.E.I.), using Filtermate 196 (Packard) cell harvester. Filters are washed with 50 mM Tris-HCl at about 0-4°C for about 4 seconds and assayed for radioactivity using Packard Top Count.

Specific binding is obtained by subtracting nonspecific binding (determined in the presence of 0.1 µM SRIF-14) from total binding. Binding

- 23 -

data are analyzed by computer-assisted nonlinear regression analysis (Data Analysis Toolbox, v.1.0, Molecular Design Limited, San Leandro, CA) and inhibition constant (K_i) values are determined.

Whether a compound of the instant invention is an SST agonist or antagonist of somatostatin is determined by the following assay.

Functional Assay: Inhibition of cAMP Intracellular Production

CHO-K1 Cells expressing human somatostatin (SRIF-14) subtype receptors are seeded in 24-well tissue culture multidishes in RPMI 1640 media with 10% fetal calf serum (FCS). The medium is changed the day before the experiment.

Cells at 10^5 cells/well are washed 2 times by 0.5 ml RPMI 1640 media. Fresh RPMI 1640 media with 0.2% BSA and supplemented with 0.5mM 3-isobutyl-1-methylxanthine ("IBMX") is added, and the cells are incubated for about 5 minutes at about 37°C. Cyclic AMP production is stimulated by the addition of 1mM forskolin ("FSK") (Sigma Chemical Co., St. Louis, MO) for about 15-30 minutes at about 37°C.

The agonist effect of a compound is measured by the simultaneous addition of FSK ($1\mu\text{M}$), SRIF-14 (Bachem, Torrence, CA), (10^{-12} M to 10^{-6} M) and a test compound (10^{-10} M to 10^{-5} M). The antagonist effect of a compound is measured by the simultaneous addition of FSK ($1\mu\text{M}$), SRIF-14 (1 to 10 nM) and a test compound (10^{-10} M to 10^{-5} M).

The reaction medium is removed and 200 ml 0.1 N HCl is added. cAMP is measured using radioimmunoassay method (Kit FlashPlate SMP001A, New England Nuclear, Boston).

Compounds of the instant invention can be and were assessed for its ability to bind to a neuromedin B receptor according to the following assay.


Cell Culture: Balb 3T3 cells, expressing the rat NMB receptor, were obtained from Dr. R.T. Jensen (National Institutes of Health, Bethesda, MD), and cultured in Dulbecco's modified Eagle's medium ("DMEM") containing

- 24 -

10% fetal calf serum, 0.5 mg/ml of G418 (Gibco). The cells were maintained at 37°C in a humidified atmosphere of 5% CO₂/95% air.

Radioligand Binding: Membranes were prepared for radioligand binding studies by homogenization of the cells in 20 ml of ice-cold 50 mM Tris-HCl with a Brinkman Polytron (Westbury, NY) (setting 6, 15 sec). The homogenates were washed twice by centrifugation (39,000g / 10 minutes), and the final pellets were resuspended in 50 mM Tris-HCl containing 5.0 mM MgCl₂, and 0.1% BSA. For assay, aliquots (0.4 ml) were incubated with 0.05 nM [¹²⁵I-Tyr⁴]bombesin (2200 Ci/mmol, New England Nuclear, Boston, MA), with and without 0.05 ml of unlabeled competing test peptides. After incubation (30 minutes, 4 °C), the bound [¹²⁵I-Tyr⁴]bombesin was separated from the free by rapid filtration through GF/C filters (Brandel, Gaithersburg, MD), which had been previously soaked in 0.3% polyethyleneimine. The filters were then washed three times with 5-ml aliquots of ice-cold 50 mM Tris-HCl, and the bound radioactivity trapped on the filters was counted by gamma spectrometry (Wallac LKB, Gaithersburg, MD). Specific binding was defined as the total [¹²⁵I-Tyr⁴]bombesin bound minus that bound in the presence of 1000 nM neuromedin B (Bachem, Torrence, CA).

One embodiment of the method includes the step of contacting the cells with a peptide of Formula (II):



or a pharmaceutically acceptable salt thereof,
wherein

AA¹ is absent or the D- or L-isomer of an amino acid selected from the group consisting of R¹¹, Aac, Aic, Arg, Asn, Asp, Dip, Gln, Glu, Hyp, Lys, Mac, Macab, Orn, Pip, Pro, Ser, Ser(Bzl), Thr, Thr(Bzl), Pip, hArg, Bip, Bpa, Tic, Cmp, , Inc, Inp, Nip, Ppc, Htic, Thi, Tra, Cmpi, Tpr, , lia, Alla, Aba, Gba, Car,

- 25 -

Ipa, Iaa, Inip, Apa, Mim, Thnc, Sala, Aala, Thza, Thia, Bal, Fala, Pala, Dap, Agly, Pgly, Ina, Dipa, Mnf, Inic, I-lqc, 3-lqc, C4c, 5-lqs, Htqa, 4-Mqc, Thn, α -Chpa, Cit, Nua, Pyp and an optionally substituted aromatic α -amino acid,

wherein said optionally substituted aromatic α -amino acid is optionally substituted with one or more substituents selected from the group consisting of halogen, NO₂, OH, CN, (C₁₋₆)alkyl, (C₂₋₆)alkenyl, (C₂₋₆)alkynyl, and NR⁹R¹⁰;

AA² is absent or the D- or L-isomer of an amino acid selected from the group consisting of R¹¹, Aic, Arg, Hca, His, Hyp, Pal, F₅-Phe, Phe, Pro, Trp, X⁰-Phe, Pip, hArg, Bip, Bpa, Tic, Cmp, , Inc, Inp, Nip, Ppc, Htic, Thi, Tra, Cmpi, Tpr, , lia, Alla, Aba, Gba, Car, Ipa, Iaa, Inip, Apa, Mim, Thnc, Sala, Aala, Thza, Thia, Bal, Fala, Pala, Dap, Agly, Pgly, Ina, Dipa, Mnf, Inic, I-lqc, 3-lqc, C4c, 5-lqs, Htqa, 4-Mqc, Thn, α -Chpa, Cit, Nua, and Pyp; AA³ is the D- or L-isomer of an amino acid selected from the group consisting of Cys, hCys, Pen, Tpa and Tmpa;

AA⁴ is a D- or L-isomer of an amino acid selected from the group consisting of Trp, N-Met-Trp, β -Met-Trp, His, hHis, hArg, Bip, Tic, , Htic, Dip, Sala, Aala, Thza, Thia, Bal, Fala, Pala, and an optionally substituted aromatic α -amino acid,

wherein said optionally substituted aromatic α -amino acid is optionally substituted with one or more substituents each independently selected from the group consisting of halogen, NO₂, OH, (C₁₋₄)alkyl, (C₂₋₄)alkenyl, (C₂₋₄)alkynyl, Bzl, O-Bzl, and NR⁹R¹⁰;

AA⁵ is a D- or L-isomer of an amino acid selected from the group consisting of 4-Pip-Gly, 4-Pip-Ala, *cis*-4-Acha, *trans*-4-Acha, *trans*-4-Amcha, hLys, Lys, Orn, hArg, Bip, Tic, , Htic, Dip, Sala, Aala, Thza, Thia, Bal, Fala, and Pala, wherein the side-chain amino group of said amino acid is optionally mono- or di-substituted with R³ and R⁴;


- 26 -

AA⁶ is a D- or L-isomer of an amino acid selected from the group consisting of Cys, hCys, Pen, Tpa, and Tmpa;

AA⁷ is absent or a D- or L-isomer of an amino acid selected from the group consisting of R¹¹, Aic, A3c, A4c, A5c, A6c, Abu, Aib, β-Ala, Arg, Bpa, Cha, Deg, Gaba, His, Ile, Leu, Nal, Nle, Pal, Phe, F₅-Phe, Pro, Sar, Ser, Ser(Bzl), Thr, Thr(Bzl), Trp, N-Me-Trp, Val, N-Me-Val, hArg, Bip, Tic, , Htic, Dip, Sala, Aala, Thza, Thia, Bal, Fala, Pala, and X⁰-Phe ;

AA⁸ is absent or the D- or L-isomer of an amino acid selected from the group consisting of R¹¹, an optionally substituted aromatic α-amino acid, Maa, Maaab, Ser, Ser(Bzl), Thr, Thr(Bzl), Tyr, Phe(4-O-Bzl), F₅-Phe, and X⁵-Phe;

R¹³ is a moiety according to the formula




wherein R²¹ is (C₁₋₄)alkyl and s is 1, 2, 3, or 4; and

X⁰ is halogen, NO₂, CH₃, OH, Bzl, O-Bzl or CN;

provided that at least one of AA⁷ or AA⁸ is present.

Another embodiment of the method includes the step of contacting the cells with a peptide of Formula (III):



or a pharmaceutically acceptable salt thereof,

wherein

AA¹ is absent or the D- or L-isomer of an amino acid selected from the group consisting of R¹¹, Aac, Aic, Arg, Asn, Asp, Gln, Glu, Hca, His, Hyp, Lys, Mac, Macab, Orn, Pro, Ser, Ser(Bzl), Thr, Thr(Bzl), Pip, hArg, Bip, Bpa, Tic, Cmp, , Inc, Inp, Nip, Ppc, Htic, Thi, Tra, Cmpi, Tpr, , lia, Alla, Aba, Gba, Car, Ipa, laa, Inip, Apa, Mim, Thnc, Sala, Aala, Thza, Thia, Bal, Fala, Pala, Dap, Agly,

- 27 -

Pgly, Ina, Dipa, Mnf, Inic, I-lqc, 3-lqc, C4c, 5-lqs, Htqa, 4-Mqc, Thn, α -Chpa, Cit, Nua, Pyp and an optionally substituted aromatic α -amino acid,

wherein said optionally substituted aromatic α -amino acid is optionally substituted with one or more substituents selected from the group consisting of halogen, NO₂, OH, CN, (C₁₋₆)alkyl, (C₂₋₆)alkenyl, (C₂₋₆)alkynyl, and NR⁹R¹⁰;

AA³ is a D- or L-isomer of an amino acid selected from the group consisting of Cys, hCys, Pen, Tpa, and Tmpa;

AA^{3b} is the D- or L-isomer of an amino acid selected from the group

consisting of R¹¹, Arg, Bpa, F₅-Phe, His, Nal, Pal, 4-Pal, Phe, Trp, hArg, Bip, Tic, , Htic, Dip, Sala, Aala, Thza, Thia, Bal, Fala, Pala, and X⁵-Phe;

AA⁴ is a D- or L-isomer of an amino acid selected from the group consisting of Trp, N-Met-Trp, β -Met-Trp, His, hHis, hArg, Bip, Tic, , Htic, Dip, Sala, Aala, Thza, Thia, Bal, Fala, Pala, and an optionally substituted aromatic α -amino

acid;

wherein said optionally substituted aromatic α -amino acid is optionally substituted with one or more substituents each independently selected from the group consisting of halogen, NO₂, OH, CN, (C₁₋₄)alkyl, (C₂₋₄)alkenyl, (C₂₋₄)alkynyl, Bzl, O-Bzl, and NR⁹R¹⁰;

AA⁵ is a D- or L-isomer of an amino acid selected from the group consisting of 4-Pip-Gly, 4-Pip-Ala, *cis*-4-Acha, *trans*-4-Acha, *trans*-4-Amcha, hLys, Lys and Orn, and, hArg, Bip, Tic, , Htic, Dip, Sala, Aala, Thza, Thia, Bal, Fala, Pala,

wherein the side-chain amino group of said amino acid is optionally mono- or di-substituted with R³ and R⁴;

AA⁶ is a D- or L-isomer of an amino acid selected from the group consisting of Cys, hCys, Pen, Tpa, and Tmpa;

AA⁷ is absent or a D- or L-isomer of an amino acid selected from the group consisting of R¹¹, Aic, A3c, A4c, A5c, A6c, Abu, Aib, β -Ala, Arg, Bpa, Cha,

- 28 -


Deg, Gaba, His, Ile, Leu, Nal, Nle, Pal, Phe, F₅-Phe, Pro, Sar, Ser, Ser(Bzl), Thr, Thr(Bzl), Trp, N-Me-Trp, Val, N-Me-Val, hArg, Bip, Tic, , Htic, Dip, Sala, Aala, Thza, Thia, Bal, Fala, Pala, and X⁰-Phe;

X⁰ is halogen, NO₂, CH₃, OH, CN, Bzl or O-Bzl;

- 5 R¹ and R² each is, independently, H, E-, E(O)₂S-, E(O)C-, EOOC-, R¹³, or absent;

R⁵ is -OR⁶ or -NR⁷R⁸;

R¹³ is a moiety of the formula



- 10 wherein R²¹ is (C₁₋₄)alkyl and s is 1, 2, 3, or 4;

provided that:

at least one of AA¹ or AA² is present;

when AA¹ is a D- or L-isomer of Pro, Hyp, Arg, Pip, hArg, Bip, Bpa, Tic, Cmp, , Inc, Inp, Nip, Ppc, Htic, Thi, Tra, Cmpi, Tpr, , lia, Alla, Aba, Gba, Car, Ipa, Iaa, Inip, Apa, Mim, Thnc, Sala, Aala, Thza, Thia, Bal, Fala, Pala, Dap, Agly, Pgly, Ina, Dipa, Mnf, Inic, I-lqc, 3-lqc, C4c, 5-lqs, Htqa, 4-Mqc, Thn, α-Chpa, Cit, Nua, Pyp or His, AA² cannot be a D- or L-isomer of Pro, Hyp, Arg, Pip, hArg, Bip, Bpa, Tic, Cmp, , Inc, Inp, Nip, Ppc, Htic, Thi, Tra, Cmpi, Tpr, , lia, Alla, Aba, Gba, Car, Ipa, Iaa, Inip, Apa, Mim, Thnc, Sala, Aala, Thza, Thia, Bal, Fala, Pala, Dap, Agly, Pgly, Ina, Dipa, Mnf, Inic, I-lqc, 3-lqc, C4c, 5-lqs, Htqa, 4-Mqc, Thn, α-Chpa, Cit, Nua, Pyp or His;

when AA⁷ is a D- or L-isomer of Thr or of Ser, AA⁸ cannot be a D- or L-isomer of Thr or of Ser;


- 25 at least one of AA¹, AA², AA^{3b}, AA⁷, AA^{7b}, or AA⁸ is the D- or L-isomer of R¹¹; and

when one of X² or X³ is =O or =S, the other is absent;

- 29 -

or a pharmaceutically acceptable salt thereof.

Yet another embodiment of the method includes the step of contacting the cells with a peptide of Formula (IV):



wherein

AA¹ is absent, the D- or L-isomer of an amino acid selected from the group consisting of R¹¹, Aic, Hyp, Pro, Ser, Ser(Bzl), Thr, Thr(Bzl), and an optionally substituted aromatic α -amino acid;

wherein said optionally substituted aromatic α -amino acid is optionally substituted with one or more substituents each independently selected from the group consisting of halogen, NO₂, OH, (C₁₋₆)alkyl, (C₂₋₆)alkenyl, (C₂₋₆)alkynyl, (C₁₋₆)alkoxy, Bzl, O-Bzl, and NR⁹R¹⁰;

AA² is absent or the D- or L-isomer of an amino acid selected from the group consisting of R¹¹, Arg, F₅-Phe, His, Pal, Phe, Trp, and X⁰-Phe;

AA³ is the D- or L-isomer of an optionally substituted aromatic α -amino acid, wherein said optionally substituted aromatic α -amino acid is optionally substituted with one or more substituents selected from the group consisting of halogen, NO₂, OH, (C₁₋₄)alkyl, (C₂₋₄)alkenyl, (C₂₋₄)alkynyl, Bzl, O-Bzl, and NR⁹R¹⁰;

AA⁴ is a D- or L-isomer of an optionally substituted amino acid selected from the group consisting of Lys, Orn, hLys, *cis*-4-Acha, *trans*-4-Acha, *trans*-4-Amcha, 4-Pip-Gly, and 4-Pip-Ala,

wherein the side chain amino group of said optionally substituted amino acid is optionally substituted with R³ and R⁴;

AA⁵ is absent or a D- or L-isomer of R¹¹, A3c, A4c, A5c, A6c, Abu, Aib, Aic, β -Ala, Bpa, Cha, Deg, F₅-Phe, Gaba, Ile, Leu, Nal, Nle, Pal, Phe, Pro, Sar, Ser, Ser(Bzl), Thr, Thr(Bzl), Trp, N-Me-Trp, Val, N-Me-Val, or X⁰-Phe;

- 30 -


AA⁶ is absent, the D- or L-isomer of R¹¹, an aromatic α -amino acid, F₅-Phe, Phe, Thr, Thr(Bzl), Ser, Ser(Bzl), or X⁰-Phe;

AA⁷ is absent, the D- or L-isomer of R¹¹ or the D- or L-isomer of an aromatic α -amino acid;

5 AA⁸ is a D- or L- isomer of R¹¹;

R¹ is H, E-, E(O)₂S-, E(O)C-, EOC-, or R¹³;

R¹³ is a moiety of the formula



wherein R²¹ is (C₁₋₄)alkyl and s is 1, 2, 3, or 4;

10 X⁰ in the definition of AA² and AA⁵ is halogen, NO₂, OH, (C₁₋₆)alkyl, (C₁₋₆)alkoxy, mono- or di-(C₁₋₆)alkylamino, Bzl or O-Bzl;

X⁰ in the definition of AA⁶ is halogen, NO₂, OH, (C₁₋₆)alkyl, (C₁₋₆)alkoxy, mono- or di-(C₁₋₆)alkylamino, Bzl, O-Bzl, or NR⁹R¹⁰;

provided that:

15 at least one of AA¹ or AA² is present;


when AA¹ is absent, AA² and AA⁸ together form a bond; and

at least two of AA⁵, AA⁶, and AA⁷ are present;

or a pharmaceutically acceptable salt thereof.

20 ABBREVIATIONS:


(A(z))aeg (A)aeg where the amino group of the adenine moiety is protected with carbobenzyloxy, i.e.,



(A)aeg N-(2-aminoethyl)-N-(2-adeninyl-1-oxo-ethyl)-glycine

- 31 -


(C(z))aeg (C)aeg where the amino group of the cytosine moiety is protected with carbobenzyloxy,



i.e.,

(C)aeg N-(2-aminoethyl)-N-(2-cytosinyl-1-oxo-ethyl)-glycine

5 (G(z))aeg (G)aeg where the amino group of the guanine moiety is protected with carbobenzyloxy, i.e.,



(G)aeg N-(2-aminoethyl)-N-(2-guaninyl-1-oxo-ethyl)-glycine

(T)aeg N-(2-aminoethyl)-N-(2-thiminyl-1-oxo-ethyl)-glycine

- | | | |
|----|------|--|
| 10 | A3c | 1-Amino-1-cyclopropane-1-carboxylic acid |
| | A4c | 1-Amino-1-cyclobutane-1-carboxylic acid |
| | A5c | 1-Amino-1-cyclopentane-1-carboxylic acid |
| | A6c | 1-Amino-1-cyclohexane-1-carboxylic acid |
| | Aaa | 2-Aminoanthraquinone |
| 15 | Aac | an aminoalkyl carboxylic acid of the formula $H_2N-(CH_2)_n-COOH$, wherein n is 2 – 6 |
| | Aala | Anthrylalanine |
| | Aba | N- (4-aminobenzoyl)- β -alanine |
| | Abp | 4-amino-1-benzylpiperidine |
| 20 | Abu | 2-Aminobutyric Acid |
| | Ac | acetyl, i.e., $CH_3-C(O)-$; |



- 32 -

	Ach	trans-1, 4-Diaminocyclohexane
	4-Acha	3-(4-aminocyclohexyl) alanine
	Ads	1-Amino-deoxy-D-sorbitol
	aeg	Aminoethylglycine
5	Agly	Allylglycine
	Ahep	1-Amino-4- (2-hydroxyethyl) Piperazine
	Aib	2-Aminoisobutyric Acid
	Aic	2-aminoindan-2-carboxylic acid
	5Aiq	5-Amino Isoquinoline
10	Alla	Allantoic acid
	4-Amcha	3-((4-aminomethyl)cyclohexyl)alanine
	Amp	1-Amino-4-methylpiperazine
	Apa	2,3-Diaminopropionic acid
	Api	1-(3-Aminopropyl) imidazole
15	Bal	3-Benzothienylalanine
	Bip	4,4'-Biphenylalanine
	BOC	Tertiarybutyloxycarbonyl
	Bpa	3-(4-biphenyl)-alanine
	Bzl	the benzyl radical
20	Bzop	4-Benzoylphenylalanine
	C4c	Cinnoline-4-Carboxylic acid
	Car	Carnosine
	Cbz	carbobenzyloxy radical
	Cha	3-Cyclohexylalanine
25	α -Chpa	Alpha-cyclohexylphenylacetic acid
	Cit	citrinin
	Cmp	4-Carboxymethylpiperidine
	Cmpi	4-Carboxymethylpiperazine
	Cpa	2-, 3-, or 4-chloro phenylalanine, unless otherwise indicated

- 33 -


	Dap	2, 3-Diaminopropionic acid
	Dapy	2,6-Diaminopyridine
	DCM	dichloromethane
	Deg	Diethylglycine
5	D-Ga	D-Glucosamine
	Dip	3,3-Diphenylalanine
	DiPa	3,5-Diiodo-4-Pyridone-1-acetic acid
	DIPEA	diisopropylethylamine
	DMF	dimethylformamide
10	Edp	4,4'-Ethylenedipiperidine
	Edt	4,4'-Ethylenedi-m-toluidine
	F ₅ -Phe	3-(Pentafluorophenyl)-alanine
	Fala	2-Furylalanine
	Fmoc	9-Fluorenylmethoxycarbonyl
15	Fpp	1-(4-Fluorophenyl) piperazine
	Gaba	4-Aminobutyric Acid
	Gba	4-Guanidinobenzoic acid
	HATU	O-(7-azabenzotriazolyl)-1,1,3,3-tetramethyluronium hexafluorophosphate
20	HBTU	O-(benzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate
	Hca	Hydrocinnamic acid (3-phenylpropionic acid)
	hCys	homocysteine
	Hep	1-(2-Hydroxyethyl) piperazine
25	hLys	homolysine
	HOAT	1-hydroxy-7-azabenzotriazole
	Htic	1,2,3,4-tetrahydroisoquinoline-7-hydroxy-3-carboxylic acid
	Htqa	4-Hydroxy-7-Trifluoromethyl-3-quinoline carboxylic acid
	Hyd	Hydralazine

- 34 -

	Hyp	4-Hydroxyproline
	laa	N- (3-Indolylacetyl)-L-Alanine
	lia	2-Imino-1-imidazolidine acetic acid
	lna	N- (3-Indolylacetyl)-L-Phenylalanine
5	Inc	Indoline-2-Carboxylic Acid
	Inic	Isonicotinic acid
	Inip	Isonipectic acid
	lpa	3-Indole Propionic Acid
	1-lqc	1-Isoquinolinecarboxylic acid
10	3-lqc	3- Isoquinolinecarboxylic acid
	5-lqs	5-Isoquinoline sulfonic Acid
		Lys with its ε amino group substituted with R ³ and R ⁴
	Lys(diEt)	Lys with its ε amino group disubstituted by two ethyl groups
	Lys(iPr)	Lys with its ε amino group monosubstituted by an isopropyl
15		group
	Maa	a mercaptoalkyl amine of the formula HS-(CH ₂) _n -NH ₂ , wherein n is 2-6;
	Maaab	a o-, m-, or p-(mercaptoalkyl)(aminoalkyl) benzene of the
		
		formula , wherein m and n each
20		is, independently, 0, 1, or 2.
	Mac	a mercaptoalkyl carboxylic acid of the formula HS-(CH ₂) _n -COOH, wherein n is 2 - 6;

- 35 -

Macab a o-, m-, or p-(mercaptoalkyl)(carboxyalkyl) benzene of the



formula , wherein m and n each is, independently, 0, 1, or 2.

	MBHA	4-methylbenzhydramine
5	Me-Trp	Trp with its indolyl nitrogen substituted with methyl
	Mim	Mimosine
	Mnf	5-(4-methyl-2-nitrophenyl)-2-furoic acid
	Mpip	1-Methylpiperazine
	4-Mqc	4-methoxy-2-quinolinecarboxylic acid
10	Nal	3-(2-naphthyl)-alanine, unless otherwise indicated
	Nip	Nipecotic acid
	Nle	norleucine
	Nua	Nicotinuric acid
	O-Bzl	the benzyloxy radical
15	Orn	ornithine
	Orn($\begin{array}{c} \text{R}^3 \\ \text{N} \\ \text{R}^4 \end{array}$)	Orn with its amino group substituted with R ³ and R ⁴
	Pal	3-(3-Pyridyl)-alanine, unless otherwise indicated
	2-Pala	2-Pyridylalanine
	3-Pala	3-Pyridylalanine
20	4-Pala	4-Pyridylalanine
	Pap	4'-Piperazinoacetophenone
	Pen	penicillamine
	Pgly	Propargylglycine
	Phg	phenylglycine
25	Pip	pipecolic acid

- 36 -

	4-Pip-Ala	3-(4-piperidyl)alanine
	4-Pip-Gly	(4-piperidyl)glycine
	Pnf	para-Nitro-phenylalanine (i.e., 4-Nitro-phenylalanine)
	Ppc	4-Phenylpiperidine-4-carboxylic Acid
5	Pyp	3-pyridine propionic acid
	Sala	Styrylalanine
	Sar	sarcosine (i.e., N-methylglycine)
	Thi	Thiaproline
	2-Thia	2-Thienylalanine
10	3-Thia	3-Thienylalanine
	Thn	1, 2, 3, 4-Tetrahydro-2-naphthoic acid
	Thnc	1,2,3,4-Tetrahydronorbarman-3-carboxylic acid
	Thza	4-Thiazolylalanine
	Tic	1,2,3,4-tetrahydro-3-isoquinolinecarboxylic acid
15	Tmpa	3-(p-thiomethylphenyl)-alanine
	Tpa	3-(p-thiophenyl)-alanine
	Tpr	Thioprolin
	Tra	Tranexamic acid
	TrPa	Tryptamine
20	X-Phe	phenylalanine with p-, o- or m-substituents X on its benzene ring, e.g., 3-(4-chlorophenyl)-alanine
	z	carbobenzyloxy

Administration of a pharmaceutically acceptable salt of a compound covered by formula (I) into a patient whose disorder arises from biochemical activity induced by NMB or somatostatin is also within the present invention.

In other words, the peptides can be provided in the form of pharmaceutically acceptable salts, e.g., acid addition salts, or metal complexes, e.g., with zinc, iron or the like. Illustrative examples of acid addition salts are those

- 37 -

with organic acids such as acetic, lactic, pantoic, maleic, citric, malic, ascorbic, succinic, benzoic, palmitic, suberic, salicylic, tartaric, methanesulfonic or toluenesulfonic acid, those with polymeric acids such as tannic acid or carboxymethyl cellulose, and those with inorganic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid or phosphoric acid.

Other features and advantages of the present invention will be apparent from the following description of the preferred embodiments, and also from the claims. It is believed that one skilled in the art can, based on the description herein, utilize the present invention to its fullest extent. The following specific embodiments are therefore to be construed as merely illustrative and not limitative of the remainder of the disclosure in any way whatsoever. All of the documents cited herein are hereby incorporated by reference

DESCRIPTION OF THE PREFERRED EMBODIMENTS

In a preferred embodiment the invention features a compound according to Formula II, wherein

AA¹ is absent, Ac-D-Phe, or the D- or L- isomer of R¹¹, Pip, Pro, or Ser, or of an aromatic α -amino acid selected from the group consisting of Cpa, Dip, Nal, Pal, and Phe;

AA² is Aic, Pal, Phe, F₅-Phe, 4-NO₂-Phe, Trp, Tyr, Phe(4-O-Bzl), or absent;

AA³ is the D- or L- isomer of an amino acid selected from the group consisting of Pen, Cys, hCys and Tmpa;

AA⁴ is the D- or L-isomer of Trp or of His;

AA⁵ is Lys, hLys, N-Me-Lys, Orn, cis-4-Acha or 4-Pip-Ala;

AA⁶ is the D- or L-isomer of an amino acid selected from the group consisting of Cys, hCys, Pen and Tmpa;

AA⁷ is A3c, A4c, A5c, A6c, Abu, Aic, β -Ala, Gaba, Nle, F₅-Phe, Phe, Pro, Sar, Ser, Thr, Thr(Bzl), Tyr, Val or absent; and

AA⁸ is R¹¹, Nal, Thr, Thr(Bzl), Tyr, Phe(4-O-Bzl), or absent;

- 38 -

or a pharmaceutically acceptable salt thereof.

In a more preferred embodiment the invention features a compound according to the immediately foregoing, wherein

5 AA¹ is absent or the D- or L- isomer of R¹¹, Pip or Pro, or of an aromatic α -amino acid selected from the group consisting of Cpa, Dip, Nal, Pal, Phe, and Ac-Phe;

AA² is Tyr, Pal, Phe, 4-NO₂-Phe, Trp, or absent;

AA³ is a D- or L-isomer of Cys or Pen;

AA⁴ is D-Trp;

10 AA⁵ is Lys, Orn, or cis-4-Acha;

AA⁶ is a D- or L-isomer of Cys or Pen;

AA⁷ is A3c, A4c, A5c, A6c, Abu, Aic, β -Ala, Gaba, Nle, Phe, Pro, Sar, Thr, Thr(Bzl), Tyr, Val, or absent; and

AA⁸ is R¹¹, Thr, Tyr, Nal, or absent;

15 or a pharmaceutically acceptable salt thereof.

In another preferred embodiment the invention features a compound according to Formula III, wherein

20 AA¹ is R¹¹, Aic, Hca, Pro, Ser, Ser(Bzl), Trp, Tyr, or a D- or L-isomer of an aromatic α -amino acid selected from the group consisting of Cpa, Nal, Ac-Nal, Phe, Ac-Phe, 4-NO₂-Phe, and Ac-4-NO₂-Phe;

AA² is Pal, Phe, F₅-Phe, Tyr, or absent;

AA³ is a D- or L-isomer of Cys, hCys, Pen or Tmpa;

AA^{3b} is Pal, 4-Pal, His, Trp, Tyr, Phe(4-O-Bzl), Phe, or R¹¹;

AA⁴ is a D- or L-isomer of Trp or His;

25 AA⁵ is Lys, N-Me-Lys, Orn, hLys, cis-4-Acha, or 4-Pip-Ala;

AA⁶ is a D- or L-isomer of Cys, hCys, Pen or Tmpa;

AA⁷ is R¹¹, A4c, A5c, Abu, β -Ala, Gaba, Phe, F₅-Phe, Ser(Bzl), Thr, Thr(Bzl), Phe(4-O-Bzl); or absent;

- 39 -

AA^{7b} is R¹¹, Nal, F₅-Phe, X⁰-Phe or absent, wherein X⁰ is halogen, NO₂, CH₃, OH, Bzl or O-Bzl; and

AA⁸ is R¹¹, Nal, Tyr, Phe(4-O-Bzl), or absent;
or a pharmaceutically acceptable salt thereof.

5 In a more preferred embodiment the invention features a compound according to the immediately foregoing, wherein

AA¹ is R¹¹, Aic, Hca, Pro, Ser(Bzl), or a D- or L-isomer of an aromatic α -amino acid selected from the group consisting of Cpa, Nal, Ac-Nal, Phe, Ac-Phe, 4-NO₂-Phe, and Ac-4-NO₂-Phe;

10 AA² is Pal, Tyr, or absent;

AA³ is a D- or L-isomer of Cys or Pen;

AA^{3b} is R¹¹, Pal, 4-Pal, Trp, Tyr, Phe(4-O-Bzl), or Phe, wherein R¹¹ is (T)aeg;

AA⁴ is D-Trp;

AA⁵ is Lys, N-Me-Lys, Orn, or cis-4-Acha;

15 AA⁶ is a D- or L-isomer of Cys or Pen;

AA⁷ is R¹¹, A5c, Abu, Ser(Bzl), Thr, Thr(Bzl), Phe(4-O-Bzl), Gaba, or absent;

AA^{7b} is Nal, X⁰-Phe or absent; and

AA⁸ is Tyr or absent;

or a pharmaceutically acceptable salt thereof.

20 In yet another preferred embodiment the invention features a compound according to Formula IV, wherein

AA¹ is Aic, Hyp, Cpa, D-Cpa, Nal, Pal, Phe, Pro, R¹¹, Tyr or absent;

AA² is Phe, Trp, F₅-Phe, His, Tyr, Phe(4-O-Bzl), or R¹¹;

AA³ is a D-isomer of Trp, His, or Pal;

25 AA⁴ is Lys, N-Me-Lys, Orn, hLys, cis-4-Acha, or 4-Pip-Ala;

AA⁵ is Pal, Phe(4-O-Bzl), Thr(Bzl), Thr, Sar, Gaba, β -Ala, A4c, A5c, A6c, Abu, Aic or absent;

AA⁶ is Thr, Tyr, Ser, F₅-Phe, Cpa, Nal, or D- or L-Phe;

AA⁷ is Nal, Pal, or absent; and

- 40 -

AA⁸ is R¹¹;

or a pharmaceutically acceptable salt thereof.

In yet another more preferred embodiment the invention features a compound according to the immediately foregoing, wherein

5 AA¹ is Cpa, Nal, Pal, Phe, Tyr or absent;

AA² is Phe, Tyr, Trp, or R¹¹;

AA³ is D-Trp;

AA⁴ is Lys, N-Me-Lys, or cis-4-Acha;

AA⁵ is Pal, Phe(4-O-Bzl), Aic, Gaba, A5c or absent;

10 AA⁶ is Thr, Nal, or D- or L-Phe;

AA⁷ is absent; and

AA⁸ is R¹¹;

or a pharmaceutically acceptable salt thereof.

In still yet another preferred embodiment the invention features a compound according to Formula II, wherein R¹ and R⁵ are absent and the N-terminal amino acid and the C-terminal amino acid together form an amide bond; or a pharmaceutically acceptable salt thereof.

In still yet another preferred embodiment the invention features a compound according to Formula III, wherein R¹ and R⁵ are absent and the N-terminal amino acid and the C-terminal amino acid together form an amide bond; or a pharmaceutically acceptable salt thereof.

In a most preferred embodiment the invention features a compound according to Formula II, wherein said compound is of the formula:

25 Ac-D-Phe-Tyr-cyclo(D-Cys-D-Trp-Lys-Cys)-Abu-Thr-NH₂;
 Nal-Tyr-cyclo(Cys-D-Trp-Lys-D-Cys)-Val-Nal-NH₂;
 Nal-Tyr-cyclo(Cys-D-Trp-Lys-D-Cys)-Abu-Nal-NH₂;
 D-Dip-Tyr-cyclo(Cys-D-Trp-Lys-D-Cys)-Abu-Nal-NH₂;
 Dip-Tyr-cyclo(D-Cys-D-Trp-Lys-D-Cys)-Abu-Nal-NH₂;
 Nal-Tyr-cyclo(D-Cys-D-Trp-Lys-D-Cys)-Abu-Nal-NH₂;

- 41 -

- Dip-Tyr-cyclo(D-Cys-D-Trp-Lys-D-Cys)-Val-Nal-NH₂;
 Nal-Tyr-cyclo(D-Cys-D-Trp-Lys-D-Cys)-Val-Nal-NH₂;
 cyclo(D-Phe-Tyr-cyclo(D-Cys-D-Trp-Lys-Cys)-Abu-Thr);
 Cpa-Pal-cyclo(D-Cys-D-Trp-Lys-D-Cys)-A3c-Nal-NH₂;
 5 Cpa-Pal-cyclo(D-Cys-D-Trp-Lys-D-Cys)-A5c-Nal-NH₂;
 Cpa-Pal-cyclo(D-Cys-D-Trp-Lys-D-Cys)-A6c-Nal-NH₂;
 (G(z))aeg-cyclo(D-Cys-D-Trp-Lys-D-Cys)-A5c-Nal-NH₂;
 Pal-cyclo(D-Cys-D-Trp-Lys-D-Cys)-A5c-Nal-NH₂;
 Cpa-Pal-cyclo(D-Cys-D-Trp-Lys-D-Cys)-β-Ala-Nal-NH₂;
 10 Cpa-Pal-cyclo(D-Cys-D-Trp-Lys-D-Cys)-Sar-Nal-NH₂;
 Cpa-Pal-cyclo(D-Cys-D-Trp-Lys-D-Cys)-Gaba-Nal-NH₂; or
 Cpa-Pal-cyclo(D-Cys-D-Trp-Lys-D-Cys)-Pro-Nal-NH₂;

or a pharmaceutically acceptable salt thereof.


- In another most preferred embodiment the invention features a
 15 compound according to Formula II, wherein said compound is of the formula:

- Phe-cyclo(Cys-D-Trp-Lys-Cys)-Thr-NH₂;
 Phe-Tyr-cyclo(D-Cys-D-Trp-Lys-Cys)-Abu-Thr-NH₂;
 Ac-D-Phe-Tyr-cyclo(D-Cys-D-Trp-Lys-Cys)-Abu-Thr-NH₂;
 Nal-Tyr-cyclo(Cys-D-Trp-Lys-D-Cys)-Val-Nal-NH₂;
 20 Nal-Tyr-cyclo(Cys-D-Trp-Lys-D-Cys)-Abu-Nal-NH₂;
 Dip-Tyr-cyclo(D-Cys-D-Trp-Lys-D-Cys)-Abu-Nal-NH₂;
 Nal-Tyr-cyclo(D-Cys-D-Trp-Lys-D-Cys)-Abu-Nal-NH₂;
 Dip-Tyr-cyclo(D-Cys-D-Trp-Lys-D-Cys)-Val-Nal-NH₂;
 Nal-Tyr-cyclo(D-Cys-D-Trp-Lys-D-Cys)-Val-Nal-NH₂;
 25 Cpa-Pal-cyclo(D-Cys-D-Trp-Lys-D-Cys)-A3c-Nal-NH₂;
 Cpa-Pal-cyclo(D-Cys-D-Trp-Lys-D-Cys)-A5c-Nal-NH₂;
 Cpa-Pal-cyclo(D-Cys-D-Trp-Lys-D-Cys)-A6c-Nal-NH₂;
 (G(z))aeg-cyclo(D-Cys-D-Trp-Lys-D-Cys)-A5c-Nal-NH₂;
 D-Cpa-cyclo(Cys-D-Trp-Lys-D-Cys)-A5c-Nal-NH₂;

- 42 -

- Pal-cyclo(D-Cys-D-Trp-Lys-D-Cys)-A5c-Nal-NH₂;
 Cpa-cyclo(D-Cys-D-Trp-Lys-D-Cys)-A5c-Nal-NH₂;
 Cpa-Pal-cyclo(D-Cys-D-Trp-Lys-D-Cys)-β-Aia-Nal-NH₂;
 Cpa-Pal-cyclo(D-Cys-D-Trp-Lys-D-Cys)-Sar-Nal-NH₂;
 5 Cpa-Pal-cyclo(D-Cys-D-Trp-Lys-D-Cys)-Aic-Nal-NH₂;
 Cpa-Pal-cyclo(D-Cys-D-Trp-Lys-D-Cys)-Gaba-Nal-NH₂;
 Cpa-Pal-cyclo(D-Cys-D-Trp-Lys-D-Cys)-Pro-Nal-NH₂;
 (T)aeg-cyclo(D-Cys-D-Trp-Lys-D-Cys)-(A)aeg-NH₂;
 Cpa-Pal-cyclo(D-Cys-D-Trp-Lys-D-Cys)-A4c-Nal-NH₂;
 10 Cpa-Pal-cyclo(D-Cys-D-Trp-Lys-D-Cys)-Nal-NH₂;
 Pal-cyclo(D-Cys-D-Trp-Lys-D-Cys)-Nal-NH₂;
 Pro-Phe-cyclo(Cys-D-Trp-Lys-D-Cys)-Val-NH₂;
 Pro-Phe-cyclo(D-Cys-D-Trp-Lys-Cys)-Val-NH₂;
 Pip-4-NO₂-Phe-cyclo(D-Cys-D-Trp-Lys-D-Cys)-Nle-NH₂;
 15 (G)aeg-Pal-cyclo(D-Cys-D-Trp-Lys-D-Cys)-Thr(Bzl)-(C)aeg-NH₂;
 (C)aeg-Pal-cyclo(D-Cys-D-Trp-Lys-D-Cys)-Thr(Bzl)-(G)aeg-NH₂;
 Pro-Phe-c(D-Cys-D-Trp-Lys-D-Cys)-Nle-Phe-NH₂;
 Pro-Phe-c(D-Cys-D-Trp-Lys-D-Cys)-Thr-Nle-NH₂;
 Pro-Phe-c(D-Cys-D-Trp-Lys-D-Cys)-Thr-Phe-NH₂;
 20 Cpa-Phe-c(D-Cys-D-Trp-Lys-D-Cys)-Gaba-NH₂;
 Cpa-Phe-c(D-Cys-D-Trp-Lys-D-Cys)-Gaba-Tyr-NH₂;
 Pip-Phe-c(D-Cys-D-Trp-Lys-D-Cys)-NH₂;
 Pip-Phe-c(Cys-D-Trp-Lys-Cys)-Gaba-NH₂; or
 Pro-Phe-c(D-Cys-D-Trp-Lys-D-Cys)-Thr-NH₂;
 25 or a pharmaceutically acceptable salt thereof.

In yet another most preferred embodiment the invention features a compound according to Formula III, wherein said compound is of the formula:



- 43 -


- D-Nal-cyclo(D-Cys-Tyr-D-Trp-Lys-Cys)-Nal-NH₂;
D-Phe-cyclo(Cys-Tyr-D-Trp-Lys-Cys)-Thr-NH₂;
D-4-NO₂-Phe-cyclo(D-Cys-Tyr-D-Trp-Lys-Cys)-Nal-NH₂;
Ac-D-4-NO₂-Phe-cyclo(D-Cys-Tyr-D-Trp-Lys-Cys)-Nal-NH₂;
5 D-4-NO₂-Phe-Pal-cyclo(D-Cys-Phe(4-O-Bzl)-D-Trp-Lys-Cys)-Tyr-NH₂;
Cpa-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;
D-4-NO₂-Phe-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Thr-Tyr-NH₂;
D-4-NO₂-Phe-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Thr(Bzl)-NH₂;
D-4-NO₂-Phe-cyclo(D-Cys-Pal-D-Trp-Lys-D-Cys)-Thr(Bzl)-Tyr-NH₂;
10 D-4-NO₂-Phe-cyclo(D-Cys-Tyr-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;
4-NO₂-Phe-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;
D-Nal-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;
Pro-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;
Cpa-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Thr(Bzl)-Nal-NH₂;
15 Ser(Bzl)-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Thr-Tyr-NH₂;
(T)aeg-cyclo(D-Cys-Pal-D-Trp-Lys-D-Cys)-Thr(Bzl)-Tyr-NH₂;
(A)aeg-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;
(G)aeg-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;
(T)aeg-cyclo(D-Cys-4-Pal-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;
20 (T)aeg-cyclo(D-Cys-Tyr-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;
(T)aeg-cyclo(D-Cys-Phe-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;
(T)aeg-cyclo(D-Cys-(T)aeg-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;
(T)aeg-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Ser(Bzl)-Tyr-NH₂;
(T)aeg-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Phe(4-O-Bzl)-Tyr-NH₂;
25 (T)aeg-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-A5c-Tyr-NH₂;
(T)aeg-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Abu-Tyr-NH₂;
D-Cpa-cyclo(D-Cys-(T)aeg-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;
(C)aeg-c(D-Cys-Pal-D-Trp-Lys-D-Cys)-Thr(Bzl)-Tyr-NH₂;
D-Cpa-c(D-Cys-Pal-D-Trp-Lys-D-Cys)Thr(Bzl)-Tyr-NH₂;

- 44 -

- (T)aeg-c(Pen-Pal-D-Trp-Lys-D-Cys)Thr(Bzl)-Tyr-NH₂;
 (T)aeg-c(D-Cys-Trp-D-Trp-Lys-D-Cys)Thr(Bzl)-Tyr-NH₂;
 (T)aeg-c(D-Cys-Phe-D-Trp-Lys-D-Cys)Thr(Bzl)-Tyr-NH₂;
 (T)aeg-c(D-Cys-Pal-D-Trp-Orn-D-Cys)Thr(Bzl)-Tyr-NH₂;
 5 (T)aeg-c(D-Cys-Pal-D-Trp-hLys-D-Cys)Thr(Bzl)-Tyr-NH₂;
 (T)aeg-c(D-Cys-Pal-D-Trp-Iamp-D-Cys)Thr(Bzl)-Tyr-NH₂;
 (T)aeg-c(D-Cys-Pal-D-Trp-Cha(4-am)-D-Cys)Thr(Bzl)-Tyr-NH₂;
 (T)aeg-c(D-Cys-Pal-D-Trp-Lys-D-Cys)-Ser(Bzl)-Tyr-NH₂;
 (T)aeg-c(D-Cys-Pal-D-Trp-Lys-D-Cys)Thr(Bzl)-D-Tyr-NH₂;
 10 (T)aeg-c(D-Cys-Pal-D-Trp-Lys-D-Cys)Thr(Bzl)-Trp-NH₂;
 (T)aeg-c(D-Cys-Pal-D-Trp-Lys-D-Pen)Thr(Bzl)-Tyr-NH₂;
 (C)aeg-c(D-Cys-Phe-D-Trp-Lys-D-Cys)Thr(Bzl)-Tyr-NH₂;
 Ina-c(D-Cys-Phe-D-Trp-Lys-D-Cys)-Thr(Bzl)-Tyr-NH₂;
 Mnf-c(D-Cys-Phe-D-Trp-Lys-D-Cys)-Thr(Bzl)-Tyr-NH₂;
 15 Inp-c(D-Cys-Phe-D-Trp-Lys-D-Cys)-Thr(Bzl)-Tyr-NH₂;
 Nua-c(D-Cys-Phe-D-Trp-Lys-D-Cys)-Thr(Bzl)-Tyr-NH₂;
 Pyp-c(D-Cys-Phe-D-Trp-Lys-D-Cys)-Thr(Bzl)-Tyr-NH₂;
 c(D-Cys-Phe-D-Trp-Lys-D-Cys)-Thr(Bzl)-Tyr-NH₂;
 (T)aeg-Pal-c(D-Cys-D-Trp-Lys-D-Cys)Thr(Bzl)-Tyr-NH₂;
 20 (T)aeg-Pal-c(D-Cys-D-Trp-Lys-D-Cys)Tyr(Bzl)-Thr-NH₂;
 (C)aeg-Phe-c(D-Cys-D-Trp-Lys-D-Cys)Thr(Bzl)-Tyr-NH₂;
 (T)aeg-D-Trp-c(D-Cys-Pal-Lys-D-Cys)Thr(Bzl)-Leu-NH₂;

or a pharmaceutically acceptable salt thereof.

- In still yet another most preferred embodiment the invention features
 25 a compound according to Formula III, wherein said compound is of the
 formula:



- 45 -

- Ac-D-Nal-cyclo(D-Cys-Tyr-D-Trp-Lys-Cys)-Nal-NH₂;
 D-Phe-cyclo(D-Cys-Tyr-D-Trp-Lys-Cys)-Nal-NH₂;
 Nal-cyclo(D-Cys-Tyr-D-Trp-Lys-Cys)-Nal-NH₂;
 D-Nal-cyclo(D-Cys-Tyr-D-Trp-Lys-Cys)-Nal-NH₂;
 5 D-Phe-cyclo(Cys-Tyr-D-Trp-Lys-Cys)-Thr-NH₂;
 D-4-NO₂-Phe-cyclo(D-Cys-Tyr-D-Trp-Lys-Cys)-Nal-NH₂;
 Ac-D-4-NO₂-Phe-cyclo(D-Cys-Tyr-D-Trp-Lys-Cys)-Nal-NH₂;
 D-4-NO₂-Phe-Pal-cyclo(D-Cys-Phe(4-O-Bzl)-D-Trp-Lys-Cys)-Tyr-NH₂;
 D-4-NO₂-Phe-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;
 10 Cpa-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;
 D-4-NO₂-Phe-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Thr(Bzl)-NH₂;
 D-4-NO₂-Phe-cyclo(D-Cys-Pal-D-Trp-Lys-D-Cys)-Thr(Bzl)-Tyr-NH₂;
 D-4-NO₂-Phe-cyclo(D-Cys-Tyr-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;
 4-NO₂-Phe-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;
 15 D-Nal-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;
 Pro-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;
 Cpa-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Thr(Bzl)-Nal-NH₂;
 Ser(Bzl)-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Thr-Tyr-NH₂;
 (T)aeg-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;
 20 (C)aeg-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;
 Aic-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;
 (C(z))aeg-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;
 (A(z))aeg-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;
 (T)aeg-cyclo(D-Cys-Pal-D-Trp-Lys-D-Cys)-Thr(Bzl)-Tyr-NH₂;
 25 (A)aeg-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;
 (G)aeg-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;
 (T)aeg-cyclo(D-Cys-4-Pal-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;
 (T)aeg-cyclo(D-Cys-Tyr-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;
 (T)aeg-cyclo(D-Cys-Phe-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;

- 46 -

- (T)aeg-cyclo(D-Cys-(T)aeg-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;
 (T)aeg-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Ser(Bzl)-Tyr-NH₂;
 (T)aeg-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Phe(4-O-Bzl)-Tyr-NH₂;
 (T)aeg-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-A5c-Tyr-NH₂;
 5 (T)aeg-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Abu-Tyr-NH₂;
 D-Cpa-cyclo(D-Cys-(T)aeg-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;
 (T)aeg-cyclo(D-Cys-Pal-D-Trp-Lys-D-Cys)-Thr(Bzl)-p-Me-Phe-NH₂;
 Ac-(T)aeg-cyclo(D-Cys-Pal-D-Trp-Lys-D-Cys)-Thr(Bzl)-Tyr-NH₂;
 (T)aeg-cyclo(D-Cys-Pal-D-Trp-Lys-D-Cys)-Nal-NH₂;
 10 D-Cpa-cyclo(D-Cys-Pal-D-Trp-Lys-D-Cys)-Nal-NH₂;
 (A)aeg-cyclo(D-Cys-Pal-D-Trp-Lys-D-Cys)-Thr(Bzl)-Tyr-NH₂; or
 (C)aeg-cyclo(D-Cys-Pal-D-Trp-Lys-D-Cys)-Thr(Bzl)-Tyr-NH₂;

or a pharmaceutically acceptable salt thereof.

- In still another most preferred embodiment the invention features a
 15 compound according to Formula IV, wherein said compound is of the
 formula:

- cyclo(Trp-D-Trp-Lys-Phe(4-O-Bzl)-Phe-(T)aeg);
 cyclo(Trp-D-Trp-Lys-Pal-Phe -(T)aeg); or
 cyclo(Phe-Phe-D-Trp-Lys-Thr-(T)aeg);
 20 or a pharmaceutically acceptable salt thereof.


PREPARATION OF PEPTIDES

- Peptides were synthesized on Rink Amide MBHA resin, (4-(2', 4'-
 dimethoxyphenyl-Fmoc-aminomethyl)-phenoxyacetamido-norleucyl-MBHA
 25 resin), using a standard solid phase protocol of Fmoc chemistry and
 cleaved with a TFA/Phenol/H₂O/triisopropylsilane (83ml/5g/10ml/2ml) mixture.
 Peptides were cyclized in CH₃CN/H₂O (5ml/5ml) using EKATHIOX™ resin
 (EKAGEN Corporation, San Carlos, CA) and purified on C₁₈ silica (Rainin
 Instruments Co., Woburn, MA, now Varian Analytical, Walnut Creek, CA),


- 47 -

using acetonitrile/0.1% trifluoroacetic acid buffers. Homogeneity was assessed by analytical HPLC and mass spectrometry and was determined to be >95% for each peptide.

Peptides having general structure



or



10 that is, having a cyclic tetra- or pentapeptyl backbone, were synthesized on Rink Amide MBHA resin, (4-(2',4'-dimethoxyphenyl-Fmoc-aminomethyl)-phenoxyacetamidonor-leucyl-MBHA resin), following a standard solid phase protocol of Fmoc-chemistry until the desired peptide was assembled. Final cleavage/deprotection was achieved by the treatment of the peptide-resin
15 with a cocktail of TFA/Phenole/H₂O/Triisopropylsilane (83:5:10:2 mL/g/mL/mL).


Cyclization (S-S bond formation) was achieved by dissolving the linear peptide in a 50% mixture of CH₃CN/H₂O, except where otherwise indicated, followed by the addition of 2.5 eq. of EKATHIOX resin then stirring
20 overnight.

Peptides were purified on C₁₈ silica column using acetonitrile/0.1%TFA buffer. Homogeneity was assessed by analytical HPLC and MAS spectrometry and was determined to be >95% for each peptide except where otherwise indicated.

25 Peptides having a carboxylic function at their C-Terminal were synthesized on Wang resin (p-Benzyloxybenzyl Alcohol resin), cleaved from

resin and deprotected by cocktail B (TFA:Phenole:H₂O:Triisopropylsilane in the ratio 88:5:5:2).

Head-To-Tail cyclic peptides having the general structure of



(IV)

were synthesized first as a totally protected linear peptide on 2-chlorotriyl chloride resin.

The first Fmoc deprotection was carried out using 5% piperidine in DMF/DCM (1:1) for about 10 minutes followed by 25% piperidine in DMF for about 15 minutes. All subsequent deprotections were performed using a standard solid phase protocol of Fmoc chemistry.




Protected linear peptides were obtained by treating the resin with acetic acid/TFE/DCM (1:1:8 by vol.) for about 60 minutes at room temperature.

Head-to-tail cyclization was achieved using HATU/HOAT/DIPEA as the coupling/cyclization reagent. The all-protected cyclic peptide was treated with a cocktail of TFA/Phenole/H₂O/Triisopropylsilane (83:5:10:2 mL/g/mL/mL) for about 2 1/2 hours to achieve final deprotection.

Peptides were purified on C₁₈ silica column using acetonitrile/0.1%TFA buffer as eluant. Homogeneity was assessed by analytical HPLC and MAS spectrometry and was determined to be >97% pure for each peptide.

As noted above, certain compounds of the invention incorporate one or more of the amino acid moiety R¹¹, having the structure

- 49 -



- wherein R^{12} , X^1 , X^2 , X^3 , X^4 , m , n , and p each is as defined in the claims. It will be apparent to one skilled in the art of chemical synthesis that the various R^{11} amino acids may be readily synthesized using appropriate starting materials and known synthesis procedures. Examples of pertinent procedures may be found in the following publications, hereby incorporated by reference: aminoethylglycine: Tetrahedron, vol. 51, pp. 6179 (1995); Bioorganic & Medicinal Chemistry Letters, vol 5, No. 11, p.1159 (1995); Tetrahedron, vol. 53, no. 43, p. 14671 (1997); Nucleosides, Nucleotides, vol. 16 (10 & 11), p. 1893 (1997); α,α -dialkylated amino acid with nucleobase side chain, Proc. Natl. Acad. Sci. USA, vo. 92, p. 12013 (1995); aminocyclohexylglycine, Chem. Eur. J. vol. 3. No. 6, p. 912 (1997); σ -N-Boc- α -N-(thymine-1-ylacetyl)ornithine, Bioorganic & Medicinal Chemistry Letters, vol. 6, no. 7, p. 793 (1996); substituted proline, J. Chem. Soc. Perkin. Trans., vol 1, pp. 539, 547, 555 (1997); N-(aminomethyl)- β -alanine,

- 50 -

Tetrahedron Lett., vol. 36, No. 38, p. 6941 (1995); substituted ornithine, Nucleosides & Nucleotides, vol 17 (1-3), pp. 219, 339 (1998); structure vi., Tetrahedron Lett., Vol. 36, no 10, p. 1713 (1995); Tetrahedron Lett, Vol. 38, no 48, p. 8363 (1997); structure v., Tetrahedron Lett., Vol 39, p. 4707 (1998); compound iv., J. Amer. Chem. Soc., vol. 119, p. 11116 (1997); aminoproline, Bioorganic & Medicinal Chemistry Lett., vol. 7, no. 6, p. 681 (1997); chiral polynucleic acid, Tetrahedron Lett., vol. 35, no. 29, p. 5173 (1994); Bioorganic & Medicinal Chemistry Lett., vol. 4, no. 8, p. 1077 (1994).

Below is a detailed description regarding the synthesis of Analog #1.

10 Other peptides of the invention can be prepared by making appropriate modifications, within the ability of someone of ordinary skill in the art of peptide synthesis.

Step 1: Preparation of Fmoc-Nal-O-tert-Butyl-Tyr-S-trityl-D-Cys-N-in-t-Boc-D-Trp-N- ϵ -t-Boc-Lys-S-trityl-D-Cys-Abu-Nal-4-(2',4'-
15 Dimethoxyphenylamino methyl) phenoxyacetamido-norluacyl-4-methylbenzhydrylamine resin.

Rink amide MBHA resin (Novabiochem, Inc., San Diego, CA), 1 g, (0.53 mmole), was placed in reaction vessel #1 (RV-1) of a Model 90 peptide synthesizer, (Advanced ChemTech, Louisville, KY). The peptide synthesizer
20 was programmed to perform the following reaction cycle:

- a. Dimethylformamide;
- b. 25% piperidine in dimethylformamide (2 times for 15 minutes each, with 1 time wash with DMF in between);
- c. DMF washes (3 x 10 mL, 1 minute each);

25 The resin was stirred with Fmoc-Nal (2.12 mmol), 2-(1H-benzotriazol-1-yl)-1, 1, 3, 3-tetramethyluronium hexafluorophosphate (HBUT) (2.01 mmole), and diisopropylethyl amino (4.24 mmole) in dimethylformamide for about 1½ hours and the resulting amino acid resin was then cycled through steps (a) to (c) in the above wash program. The

- 51 -

Nal-resin was coupled with Fmoc-Abu, then cycled as described above. It was dried under vacuum.

The following amino acids (1.4 mmole) were coupled successively to the peptide resin (0.35 mmole), by the same procedure: Fmoc-S-Trityl-D-Cys, Fmoc-N- ϵ -t-Boc-Lys, Fmoc-N-in-t-Boc-D-Trp. The peptide resin, after drying under vacuum, was split and one portion coupled with Fmoc-S-Trityl-D-Cys, Fmoc-O-t-butyl-Tyr. The coupled portion was split again and one portion coupled with Fmoc-Nal. After washing with DMF (3 x 10 mL, 1 minute each) and drying under vacuum, the completed resin weighed 0.242 g.

Step 2: Preparation of H-Nal-Tyr-D-Cys-D-Trp-Lys-D-Cys-Abu-Nal-NH₂

The peptide resin obtained from Step 1 (0.24 g, 0.087 mmole) was mixed with a freshly prepared solution of TFA (8.8 mL), phenol (0.5g), H₂O (0.5 mL) and triisopropylsilane (0.2mL) at room temperature and stirred for about 2½ hours. Excess TFA was evaporated under reduced pressure to an oily residue. Ether was then added to the oily residue and the free linear peptide was precipitated, filtered, then washed with dry ether. The crude peptide was then dissolved in 11mL of CH₃CN/H₂O/0.1N HOAc (5mL/5mL/1mL), followed by the addition of 200 mg EKATHIOX® resin. The mixture was stirred overnight and then filtered. The filtrate was evaporated to a small volume then applied to a column (22-250 mm) of microsorb octadecylsilane silica (5µm), followed by elution with a linear gradient (30% to 80%, 30 minutes) of acetonitrile in water, in which both solvents have 0.1% trifluoroacetic acid. Fractions were examined by analytical high performance liquid chromatography ("HPLC") and pooled to give maximum purity. Lyophilization of the solutions from water gave 10 mg of the product as a white, fluffy powder. The product was found to be homogeneous by HPLC C₁₈ silica using the same eluant as immediately above, (tR = 16.646

- 52 -

minutes). Infusion mass spectrometry confirmed the composition of the cyclic octapeptide; (MW 1178.45).


OTHER EMBODIMENTS

From the above description, one skilled in the art can easily ascertain
5 the essential characteristics of the present invention, and without departing
from the spirit and scope thereof, can make various changes and
modifications of the invention to adapt it to various uses and conditions.
Thus, other embodiments are also within the claims.

CLAIMS

What is claimed is:

1. A compound of formula (I):



5

(I)

or a pharmaceutically acceptable salt thereof,

wherein

the α -nitrogen of AA^1 , AA^2 , AA^3 , AA^{3b} , AA^4 , AA^5 , AA^6 , AA^7 , AA^{7b} , and AA^8 each is, independently, optionally substituted with (C_{1-4}) alkyl, (C_{3-4}) alkenyl, (C_{3-4}) alkynyl, or (C_{1-6}) alkyl-C(O)-;

10

AA^1 is absent or the D- or L-isomer of an amino acid selected from the group consisting of R^{11} , Aac, Aic, Arg, Asn, Asp, Dip, Gln, Glu, Hca, Hyp, Lys, Mac, Macab, Orn, Pro, Ser, Ser(Bzl), Thr, Thr(Bzl), Pip, hArg, Bip, Bpa, Tic, Cmp, Inc, Inp, Nip, Ppc, Htic, Thi, Tra, Cmpi, Tpr, Iia, Alla, Aba, Gba, Car, Ipa, Iaa, Inip, Apa, Mim, Thnc, Sala, Aala, Thza, Thia, Bal, Fala, Pala, Dap, Agly, Pgly, Ina, Dipa, Mnf, Inic, C4c, 5-lqs, Htqa, 4-Mqc, Thn, α -Chpa, Cit, Nua, Pyp and an optionally substituted aromatic α -amino acid;

15

wherein said optionally substituted aromatic α -amino acid is optionally substituted with one or more substituents each independently selected from the group consisting of halogen, NO_2 , OH, CN, (C_{1-6}) alkyl, (C_{2-6}) alkenyl, (C_{2-6}) alkynyl, (C_{1-6}) alkoxy, Bzl, O-Bzl, and NR^9R^{10} ;

20

AA^2 is absent or the D- or L-isomer of an amino acid selected from the group consisting of R^{11} , Aic, Arg, Hca, His, Hyp, Pal, F₅-Phe, Phe, Pro, Trp, and X⁰-Phe Pip, hArg, Bip, Bpa, Tic, Cmp, , Inc, Inp, Nip, Ppc, Htic, Thi, Tra, Cmpi, Tpr, Iia, Alla, Aba, Gba, Car, Ipa, Iaa, Inip, Apa, Mim, Thnc, Sala, Aala, Thza, Thia, Bal, Fala, Pala, Dap, Agly, Pgly, Ina, Dipa, Mnf, Inic, I-lqc, 3-lqc, C4c, 5-lqs, Htqa, 4-Mqc, Thn, α -Chpa, Cit, Nua, and Pyp;

25

- 54 -

AA³ is the D- or L-isomer of an amino acid selected from the group consisting of Cys, hCys, Pen, Tpa, Tmpa, Mac, Macab, and an optionally substituted aromatic α -amino acid;

wherein said optionally substituted aromatic α -amino acid is optionally substituted with one or more substituents selected from the group consisting of halogen, NO₂, OH, CN, (C₁₋₄)alkyl, (C₂₋₄)alkenyl, (C₂₋₄)alkynyl, (C₁₋₄)alkoxy, Bzl, O-Bzl, NR⁹R¹⁰, Pip, hArg, Bip, Bpa, Tic, Cmp, , Inc, Inp, Nip, Ppc, Htic, Thi, Tra, Cmpi, Tpr, , lia, Alla, Aba, Gba, Car, Ipa, laa, Inip, Apa, Mim, Thnc, Sala, Aala, Thza, Thia, Bal, Fala, Pala, Dap, Agly, Pgly, Ina, Dipa, Mnf, Inic, I-lqc, 3-lqc, C4c, 5-lqs, Htqa, 4-Mqc, Thn, α -Chpa, Cit, Nua, and Pyp; AA^{3b} is absent or the D- or L-isomer of an amino acid selected from the group consisting of Pal, 4-Pal, His, Arg, Nal, Trp, Bpa, F₅-Phe, Phe, X⁰-Phe, R¹¹, hArg, Bip, Tic, , Htic, Dip, Sala, Aala, Thza, Thia, Bal, Fala, and Pala;

AA⁴ is a D- or L-isomer of an optionally substituted amino acid or of an optionally substituted aromatic α -amino acid;

wherein said optionally substituted amino acid is selected from the group consisting of Trp, Lys, Orn, hLys, *cis*-4-Acha, *trans*-4-Acha, *trans*-4-Amcha, 4-Pip-Gly, N-Met-Trp, β -Met-Trp, His, hHis, hArg, Bip, Tic, Htic, Dip, Sala, Aala, Thza, Thia, Bal, Fala, Pala, and 4-Pip-Ala;

wherein the side chain amino group of said optionally substituted amino acid is optionally substituted with R³ and R⁴; and

wherein said optionally substituted aromatic α -amino acid is optionally substituted with one or more substituents each independently selected from the group consisting of halogen, NO₂, OH, CN, (C₁₋₄)alkyl, (C₂₋₄)alkenyl, (C₂₋₄)alkynyl, Bzl, O-Bzl, and NR⁹R¹⁰;

AA⁵ is absent, R¹¹, Aic, A3c, A4c, A5c, A6c, Abu, Aib, β -Ala, Bpa, Cha, Deg, Gaba, Ile, Leu, Nal, Nle, Pro, Sar, Ser, Ser(Bzl), Thr, Thr(Bzl), Trp, Val, Pal, F₅-Phe, Phe, X⁰-Phe, or an optionally substituted D- or L- isomer of an amino acid selected from the group consisting of 4-Pip-Gly, 4-Pip-Ala, *cis*-4-Acha,

- 55 -

trans-4-Acha, *trans*-4-Amcha, hLys, Lys, Orn, hArg, Bip, Tic, , Htic, Dip, Sala, Aala, Thza, Thia, Bal, Fala, and Pala; wherein the side-chain amino group of said optionally substituted amino acid is optionally mono- or di-substituted with R³ and R⁴;

- 5 AA⁶ is absent or the D- or L-isomer of an amino acid selected from the group consisting of R¹¹, an optionally substituted aromatic α -amino acid, Cys, hCys, Pen, Tpa, Tmpa, Thr, Thr(Bzl), Ser, Ser(Bzl), hArg, Bip, Tic, , Htic, Dip, Sala, Aala, Thza, Thia, Bal, Fala, and Pala;

- 10 AA⁷ is absent or the D- or L-isomer of an amino acid selected from the group consisting of R¹¹, an optionally substituted aromatic α -amino acid, A3c, A4c, A5c, A6c, Abu, Aib, Aic, β -Ala, Arg, Cha, Deg, Gaba, Ile, Leu, Nle, Pip, Pro, Sar, Ser, Ser(Bzl), Thr, Thr(Bzl), Val, Tic, Htic, Sala, Aala, Thza, Thia, Bal, Fala, Pala, hArg, Bip, Bpa, Dip, Pal, Sala, and X⁰-Phe;

- 15 AA^{7b} is absent or a D- or L-isomer of an amino acid selected from the group consisting of R¹¹, Bpa, Phe, F₅-Phe, X⁰-Phe, Nal, Pro, Ser, Ser(Bzl), Thr, Thr(Bzl), Trp, hArg, Bip, Tic, Htic, Dip, Sala, Aala, Thza, Thia, Bal, Fala, and Pala;

- 20 AA⁸ is absent or the D- or L- isomer of an amino acid selected from the group consisting of R¹¹, Maa, Maaab, Thr, Thr(Bzl), Ser, Ser(Bzl), Tyr, Phe(4-O-Bzl), F₅-Phe, and X⁵-Phe, and an optionally substituted aromatic α -amino acid;

R¹ and R² each is, independently, H, E-, E(O)₂S-, E(O)C-, EOO-, R¹³, or absent;

- 25 R³ and R⁴ each is, independently, (C₁₋₁₂)alkyl, (C₂₋₁₂)alkenyl, (C₂₋₁₂)alkynyl, phenyl, naphthyl, phenyl-(C₁₋₆)alkyl, phenyl-(C₂₋₆)alkenyl, phenyl-(C₂₋₆)alkynyl, naphthyl-(C₁₋₆)alkyl, naphthyl-(C₂₋₆)alkenyl, naphthyl-(C₂₋₆)alkynyl, (cyclo(C₃₋₇)alkyl)-(C₁₋₆)alkyl, (cyclo(C₃₋₇)alkyl)-(C₂₋₆)alkenyl, (cyclo(C₃₋₇)alkyl)-(C₂₋₆)alkynyl, heterocyclyl-(C₁₋₄)alkyl, heterocyclyl-(C₂₋₄)alkenyl, heterocyclyl-(C₂₋

- 56 -




4)alkynyl, 1-adamantyl, 2-adamantyl, 9-fluorenylmethyl, dicyclopropylmethyl, dimethylcyclopropylmethyl, or benzhydryl;

R^5 is $-OR^6$, $-NR^7R^8$, or absent,

wherein each R^6 , R^7 and R^8 is, independently, H, (C_{1-12}) alkyl, (C_{2-12}) alkenyl, (C_{2-12}) alkynyl, phenyl, naphthyl, phenyl- (C_{1-6}) alkyl, phenyl- (C_{2-6}) alkenyl, phenyl- (C_{2-6}) alkynyl, naphthyl- (C_{1-6}) alkyl, naphthyl- (C_{2-6}) alkenyl, naphthyl- (C_{2-6}) alkynyl, 1-adamantyl, 2-adamantyl, 9-fluorenylmethyl, dicyclopropylmethyl, dimethylcyclopropylmethyl, or benzhydryl;

R^9 and R^{10} each is, independently, H, (C_{1-6}) alkyl, (C_{3-4}) alkenyl, (C_{3-4}) alkynyl, 1-adamantyl, or 2-adamantyl;


R^{11} is, independently for each occurrence, a D- or L-amino acid of the formula:




- 57 -

wherein m and n each is, independently, 1, 2, or 3, and p is 0, 1, or 2;

R¹² is, independently for each occurrence, an optionally substituted moiety of the formula:



R¹³ is a moiety of the formula



wherein q, r, s, and t each is, independently, 0, 1, 2, 3, 4 or 5;

R¹⁹ is absent, H, NH₂, OH, (C₁₋₆)hydroxyalkyl, N(R²⁷R²⁸), SO₃H, or an optionally substituted moiety selected from the group consisting of heterocyclyl, phenyl and naphthyl,

wherein the optionally substituted moiety defined for R¹⁹ is optionally substituted with one or more substituents selected, independently for each occurrence, from the group consisting of halogen, NO₂, OH, (C₁₋₆)alkyl, (C₂₋₆)alkenyl, (C₂₋₆)alkynyl, (C₁₋₆)alkoxy, NH₂, mono- or di-(C₁₋₆)alkylamino, Bzl, and O-Bzl;

R²⁰ is O or absent;

R²¹ is (C₁₋₆)alkyl or absent;

- 58 -

R²² is N, O, C, or CH;

R²³ is (C₁₋₆)alkyl or absent;

R²⁴ is N, CH, or C;

R²⁵ is NH, O, or absent;

5 R²⁶ is SO₂, CO, or CH;

R²⁷ and R²⁸ each is, independently, H or (C₁₋₆)alkyl;

E is, independently for each occurrence, an optionally substituted moiety selected from the group consisting of (C₁₋₁₂)alkyl, (C₂₋₁₂)alkenyl, (C₂₋₁₂)alkynyl, phenyl, naphthyl, phenyl-(C₁₋₆)alkyl, phenyl-(C₂₋₆)alkenyl, phenyl-(C₂₋₆)alkynyl, naphthyl-(C₁₋₆)alkyl, naphthyl-(C₂₋₆)alkenyl, naphthyl-(C₂₋₆)alkynyl, (cyclo(C₃₋₇alkyl)-(C₁₋₆)alkyl, (cyclo(C₃₋₇alkyl)-(C₂₋₆)alkenyl, (cyclo(C₃₋₇alkyl)-(C₂₋₆)alkynyl, heterocyclyl-(C₁₋₄)alkyl, heterocyclyl-(C₂₋₄)alkenyl, heterocyclyl-(C₂₋₄)alkynyl, 1-adamantyl, 2-adamantyl, dicyclopropylmethyl, dimethylcyclopropylmethyl, 9-fluorenylmethyl, and benzhydryl;

15 wherein the optionally substituted moiety defined for E is optionally substituted with one or more substituents each independently selected from the group consisting of halogen, OH, Bzl, O-Bzl, NO₂, CN, COOH, and SH;

20 X⁰ is halogen, NO₂, OH, (C₁₋₆)alkyl, (C₁₋₆)alkoxy, mono- or di-(C₁₋₆)alkylamino, Bzl, O-Bzl, NR⁹R¹⁰, or CN;

X¹ is H, (C₁₋₆)alkyl, (C₂₋₆)alkenyl, (C₂₋₆)alkynyl, indolyl, imidazolyl, 1-naphthyl, 3-pyridyl, optionally ring-substituted benzyl, or a moiety which corresponds to the side-chain group of Arg, Leu, Gln, Lys, Tyr, His, Thr, Trp, Phe, Val, Ala, Lys, or His;

25 wherein said optionally ring-substituted benzyl is optionally substituted with one or more substituents selected from the group consisting of halogen, OH, (C₁₋₆)alkoxy, mono- or di-(C₁₋₆)alkylamino, (C₁₋₄) alkyl, (C₂₋₄) alkenyl, (C₂₋₄) alkynyl, and NR⁹R¹⁰;

- 59 -

X^2 and X^3 each is, independently, H, halogen, OH, =O, =S, (C_{1-12}) alkyl, (C_{2-12}) alkenyl, (C_{2-12}) alkynyl, phenyl, naphthyl, phenyl- (C_{1-6}) alkyl, phenyl- (C_{2-6}) alkenyl, phenyl- (C_{2-6}) alkynyl, naphthyl- (C_{1-6}) alkyl, naphthyl- (C_{2-6}) alkenyl, naphthyl- (C_{2-6}) alkynyl, $(cyclo(C_{3-7})alkyl)-(C_{1-6})alkyl$, $(cyclo(C_{3-7})alkyl)-(C_{2-6})alkenyl$, $(cyclo(C_{3-7})alkyl)-(C_{2-6})alkynyl$, heterocyclyl- $(C_{1-4})alkyl$, heterocyclyl- $(C_{2-4})alkenyl$, heterocyclyl- $(C_{2-4})alkynyl$, 1-adamantyl, 2-adamantyl, dicyclopropylmethyl, or dimethylcyclopropyl methyl;

X^4 is H, OH, or NH_2 ; and

X^5 is halogen, NO_2 , CH_3 , OH, Bzl or O-Bzl;

10 provided that:

at least six amino acid residues are present;

when AA^3 is a D- or L-isomer of an amino acid selected from the group consisting of Cys, hCys, Pen, Tpa, or Tmpa, and AA^6 is a D- or L-isomer of an amino acid selected from the group consisting of Cys, hCys, Pen, Tpa, or Tmpa, then AA^3 and AA^6 are connected by a disulfide bond;

15 when AA^1 or AA^3 is a D- or L-isomer of an amino acid selected from the group consisting of Mac or Macab, then AA^8 is a D- or L-isomer of an amino acid selected from the group consisting of Maa and Maaab, and

when AA^8 is a D- or L-isomer of an amino acid selected from the group consisting of Maa and Maaab, then AA^1 or AA^3 is a D- or L-isomer of Mac or of Macab, and AA^1 or AA^3 is connected by a disulfide bond with AA^8 ;

20 AA^2 can be D- or L-Hca only when AA^1 is absent;

when one of R^1 or R^2 is $E(O)_2S-$, $E(O)C-$, $EOOC-$, or R^{13} , the other is H;

when R^5 is absent, then one of R^1 or R^2 is also absent, and the N-terminal amino acid and C-terminal amino acid together form an amide bond;

25 when one of X^2 or X^3 is $C=O$ or $C=S$, the other is absent; and

said compound of formula (I) is not of the formula:

D-Phe-Tyr-cyclo(D-Cys-D-Trp-Lys-Cys)-Abu-Thr- NH_2 ;

Ac-Phe-Tyr-cyclo(D-Cys-D-Trp-Lys-Cys)-Abu-Thr- NH_2 ;


- 60 -

L-4-NO₂-Phe-Tyr-cyclo(D-Cys-D-Trp-Lys-Cys)-Abu-Thr-NH₂;Ac-L-4-NO₂-Phe-Tyr-cyclo(D-Cys-D-Trp-Lys-Cys)-Abu-Thr-NH₂;Hca-Tyr-cyclo(D-Cys-D-Trp-Lys-Cys)-Abu-Thr-NH₂;D-Dip-Tyr-cyclo(Cys-D-Trp-Lys-D-Cys)-Val-Nal-NH₂;5 D-4-NO₂-Phe-Phe(4-O-Bzl)-cyclo(D-Cys-D-Trp-Lys-Cys)Cha-Nal-NH₂;

or

D-4-NO₂-Phe-cyclo(D-Cys-Phe(4-O-Bzl)-D-Trp-Lys-Cys)-Val-Tyr-NH₂.

2. A compound according to claim 1, wherein said compound is of
10 formula (II):



or a pharmaceutically acceptable salt thereof,

wherein

15 AA¹ is absent or the D- or L-isomer of an amino acid selected from the group consisting of R¹¹, Aac, Aic, Arg, Asn, Asp, Dip, Gln, Glu, Hyp, Lys, Mac, Macab, Orn, Pip, Pro, Ser, Ser(Bzl), Thr, Thr(Bzl), Pip, hArg, Bip, Bpa, Tic, Cmp, , Inc, Inp, Nip, Ppc, Htic, Thi, Tra, Cmpi, Tpr, , lia, Alla, Aba, Gba, Car, Ipa, Iaa, Inip, Apa, Mim, Thnc, Sala, Aala, Thza, Thia, Bal, Fala, Pala, Dap,
20 Agly, Pgly, Ina, Dipa, Mnf, Inic, I-lqc, 3-lqc, C4c, 5-lqs, Htqa, 4-Mqc, Thn, α-Chpa, Cit, Nua, Pyp and an optionally substituted aromatic α-amino acid,

wherein said optionally substituted aromatic α-amino acid is optionally substituted with one or more substituents selected from the group consisting of halogen, NO₂, OH, CN, (C₁₋₆)alkyl, (C₂₋₆)alkenyl, (C₂₋₆)alkynyl,

25 and NR⁹R¹⁰;

AA² is absent or the D- or L-isomer of an amino acid selected from the group consisting of R¹¹, Aic, Arg, Hca, His, Hyp, Pal, F₅-Phe, Phe, Pro, Trp, X⁰-Phe, Pip, hArg, Bip, Bpa, Tic, Cmp, , Inc, Inp, Nip, Ppc, Htic, Thi, Tra, Cmpi, Tpr, ,

- 61 -

lia, Alla, Aba, Gba, Car, Ipa, Iaa, Inip, Apa, Mim, Thnc, Sala, Aala, Thza, Thia, Bal, Fala, Pala, Dap, Agly, Pgly, Ina, Dipa, Mnf, Inic, I-lqc, 3-lqc, C4c, 5-lqs, Htqa, 4-Mqc, Thn, α -Chpa, Cit, Nua, and Pyp; AA³ is the D- or L-isomer of an amino acid selected from the group consisting of Cys, hCys,

5 Pen, Tpa and Tmpa;

AA⁴ is a D- or L-isomer of an amino acid selected from the group consisting of Trp, N-Met-Trp, β -Met-Trp, His, hHis, hArg, Bip, Tic, , Htic, Dip, Sala, Aala, Thza, Thia, Bal, Fala, Pala, and an optionally substituted aromatic α -amino acid,

10 wherein said optionally substituted aromatic α -amino acid is optionally substituted with one or more substituents each independently selected from the group consisting of halogen, NO₂, OH, (C₁₋₄)alkyl, (C₂₋₄)alkenyl, (C₂₋₄)alkynyl, Bzl, O-Bzl, and NR⁹R¹⁰;

AA⁵ is a D- or L-isomer of an amino acid selected from the group consisting
15 of 4-Pip-Gly, 4-Pip-Ala, *cis*-4-Acha, *trans*-4-Acha, *trans*-4-Amcha, hLys, Lys, Orn, hArg, Bip, Tic, , Htic, Dip, Sala, Aala, Thza, Thia, Bal, Fala, and Pala, wherein the side-chain amino group of said amino acid is optionally mono- or di-substituted with R³ and R⁴;


AA⁶ is a D- or L-isomer of an amino acid selected from the group consisting
20 of Cys, hCys, Pen, Tpa, and Tmpa;

AA⁷ is absent or a D- or L-isomer of an amino acid selected from the group consisting of R¹¹, Aic, A3c, A4c, A5c, A6c, Abu, Aib, β -Ala, Arg, Bpa, Cha, Deg, Gaba, His, Ile, Leu, Nal, Nle, Pal, Phe, F₅-Phe, Pro, Sar, Ser, Ser(Bzl), Thr, Thr(Bzl), Trp, N-Me-Trp, Val, N-Me-Val, hArg, Bip, Tic, , Htic, Dip, Sala,
25 Aala, Thza, Thia, Bal, Fala, Pala, and X⁰-Phe ;

AA⁸ is absent or the D- or L-isomer of an amino acid selected from the group consisting of R¹¹, an optionally substituted aromatic α -amino acid, Maa, Maaab, Ser, Ser(Bzl), Thr, Thr(Bzl), Tyr, Phe(4-O-Bzl), F₅-Phe, and X⁵-Phe;

R¹³ is a moiety according to the formula


- 62 -



wherein R^{21} is (C_{1-4}) alkyl and s is 1, 2, 3, or 4; and
 X^0 is halogen, NO_2 , CH_3 , OH , Bzl , O-Bzl or CN ;
 provided that at least one of AA^7 or AA^8 is present.

5

3. A compound according to claim 1, wherein said compound is of formula (III):



10 or a pharmaceutically acceptable salt thereof,
 wherein

AA^1 is absent or the D- or L-isomer of an amino acid selected from the group consisting of R^{11} , Aac, Aic, Arg, Asn, Asp, Gln, Glu, Hca, His, Hyp, Lys, Mac, Macab, Orn, Pro, Ser, Ser(Bzl), Thr, Thr(Bzl), Pip, hArg, Bip, Bpa, Tic, Cmp, , Inc, Inp, Nip, Ppc, Htic, Thi, Tra, Cmpi, Tpr, , lia, Alla, Aba, Gba, Car, Ipa, 15 , Iaa, Inip, Apa, Mim, Thnc, Sala, Aala, Thza, Thia, Bal, Fala, Pala, Dap, Agly, Pgly, Ina, Dipa, Mnf, Inic, I-lqc, 3-lqc, C4c, 5-lqs, Htqa, 4-Mqc, Thn, α -Chpa, Cit, Nua, Pyp and an optionally substituted aromatic α -amino acid,

wherein said optionally substituted aromatic α -amino acid is optionally 20 substituted with one or more substituents selected from the group consisting of halogen, NO_2 , OH , CN , (C_{1-6}) alkyl, (C_{2-6}) alkenyl, (C_{2-6}) alkynyl, and NR^9R^{10} ;

AA^3 is a D- or L-isomer of an amino acid selected from the group consisting of Cys, hCys, Pen, Tpa, and Tmpa;

- 63 -

AA^{3b} is the D- or L-isomer of an amino acid selected from the group consisting of R¹¹, Arg, Bpa, F₅-Phe, His, Nal, Pal, 4-Pal, Phe, Trp, hArg, Bip, Tic, , Htic, Dip, Sala, Aala, Thza, Thia, Bal, Fala, Pala, and X⁵-Phe;

AA⁴ is a D- or L-isomer of an amino acid selected from the group consisting
 5 of Trp, N-Met-Trp, β -Met-Trp, His, hHis, hArg, Bip, Tic, , Htic, Dip, Sala, Aala, Thza, Thia, Bal, Fala, Pala, and an optionally substituted aromatic α -amino acid;

wherein said optionally substituted aromatic α -amino acid is optionally substituted with one or more substituents each independently selected
 10 from the group consisting of halogen, NO₂, OH, CN, (C₁₋₄)alkyl, (C₂₋₄)alkenyl, (C₂₋₄)alkynyl, Bzl, O-Bzl, and NR⁹R¹⁰;

AA⁵ is a D- or L-isomer of an amino acid selected from the group consisting of 4-Pip-Gly, 4-Pip-Ala, *cis*-4-Acha, *trans*-4-Acha, *trans*-4-Amcha, hLys, Lys and Orn, and, hArg, Bip, Tic, , Htic, Dip, Sala, Aala, Thza, Thia, Bal, Fala,
 15 Pala,

wherein the side-chain amino group of said amino acid is optionally mono- or di-substituted with R³ and R⁴;

AA⁶ is a D- or L-isomer of an amino acid selected from the group consisting of Cys, hCys, Pen, Tpa, and Tmpa;

20 AA⁷ is absent or a D- or L-isomer of an amino acid selected from the group consisting of R¹¹, Aic, A3c, A4c, A5c, A6c, Abu, Aib, β -Ala, Arg, Bpa, Cha, Deg, Gaba, His, Ile, Leu, Nal, Nle, Pal, Phe, F₅-Phe, Pro, Sar, Ser, Ser(Bzl), Thr, Thr(Bzl), Trp, N-Me-Trp, Val, N-Me-Val, hArg, Bip, Tic, , Htic, Dip, Sala, Aala, Thza, Thia, Bal, Fala, Pala, and X⁰-Phe;


25 X⁰ is halogen, NO₂, CH₃, OH, CN, Bzl or O-Bzl;

R¹ and R² each is, independently, H, E-, E(O)₂S-, E(O)C-, EOOC-, R¹³, or absent;

R⁵ is -OR⁶ or -NR⁷R⁸;

R¹³ is a moiety of the formula

- 64 -




wherein R^{21} is (C_{1-4}) alkyl and s is 1, 2, 3, or 4;

provided that:

- at least one of AA^1 or AA^2 is present;
- 5 when AA^1 is a D- or L-isomer of Pro, Hyp, Arg, Pip, hArg, Bip, Bpa, Tic, Cmp, , Inc, Inp, Nip, Ppc, Htic, Thi, Tra, Cmpi, Tpr, , lia, Alla, Aba, Gba, Car, Ipa, Iaa, Inip, Apa, Mim, Thnc, Sala, Aala, Thza, Thia, Bal, Fala, Pala, Dap, Agly, Pgly, Ina, Dipa, Mnf, Inic, I-lqc, 3-lqc, C4c, 5-lqs, Htqa, 4-Mqc, Thn, α -Chpa, Cit, Nua, Pyp or His, AA^2 cannot be a D- or L-isomer
- 10 of Pro, Hyp, Arg, Pip, hArg, Bip, Bpa, Tic, Cmp, , Inc, Inp, Nip, Ppc, Htic, Thi, Tra, Cmpi, Tpr, , lia, Alla, Aba, Gba, Car, Ipa, Iaa, Inip, Apa, Mim, Thnc, Sala, Aala, Thza, Thia, Bal, Fala, Pala, Dap, Agly, Pgly, Ina, Dipa, Mnf, Inic, I-lqc, 3-lqc, C4c, 5-lqs, Htqa, 4-Mqc, Thn, α -Chpa, Cit, Nua, Pyp or His;
- 15 when AA^7 is a D- or L-isomer of Thr or of Ser, AA^8 cannot be a D- or L-isomer of Thr or of Ser;
- at least one of AA^1 , AA^2 , AA^{3b} , AA^7 , AA^{7b} , or AA^8 is the D- or L-isomer of R^{11} ; and
- when one of X^2 or X^3 is $=\text{O}$ or $=\text{S}$, the other is absent;
- 20 or a pharmaceutically acceptable salt thereof.

4. A compound according to claim 1, wherein said compound is of formula (IV):



wherein

AA¹ is absent or the D- or L-isomer of an amino acid selected from the group consisting of R¹¹, Aic, Hyp, Pro, Ser, Ser(Bzl), Thr, Thr(Bzl), Tic, Htic, Fala and an optionally substituted aromatic α -amino acid;

- 5 wherein said optionally substituted aromatic α -amino acid is optionally substituted with one or more substituents each independently selected from the group consisting of halogen, NO₂, OH, CN, (C₁₋₆)alkyl, (C₂₋₆)alkenyl, (C₂₋₆)alkynyl, (C₁₋₆)alkoxy, Bzl, O-Bzl, and NR⁹R¹⁰;

- AA² is absent or the D- or L-isomer of an amino acid selected from the group consisting of R¹¹, Arg, F₅-Phe, His, Pal, Phe, Trp, hArg, Pala, Bal, Fala, ,
10 Sala and X⁰-Phe;

- AA³ is the D- or L-isomer of an optionally substituted aromatic α -amino acid, wherein said optionally substituted aromatic α -amino acid is optionally substituted with one or more substituents selected from the group consisting of halogen, NO₂, OH, CN, (C₁₋₄)alkyl, (C₂₋₄)alkenyl, (C₂₋₄)alkynyl, Bzl, O-Bzl,
15 and NR⁹R¹⁰;

AA⁴ is a D- or L-isomer of an optionally substituted amino acid selected from the group consisting of Trp, N-Met-Trp, β -Me-Trp, Lys, Orn, hLys, *cis*-4-Acha, *trans*-4-Acha, *trans*-4-Amcha, 4-Pip-Gly, 4-Pip-Ala, hArg, Bip, Tic, Htic, Dip, Sala, Aala, Thza, Thia, Bal, Fala, and Pala;

- 20 wherein the side chain amino group of said optionally substituted amino acid is optionally substituted with R³ and R⁴;

- AA⁵ is absent or a D- or L-isomer of R¹¹, A3c, A4c, A5c, A6c, Abu, Aib, Aic, β -Ala, Bpa, Cha, Deg, F₅-Phe, Gaba, Ile, Leu, Nal, Nle, Pal, Phe, Pro, Sar, Ser, Ser(Bzl), Thr, Thr(Bzl), Trp, N-Me-Trp, Val, N-Me-Val, hArg, Bip, Tic, ,
25 Htic, Dip, Sala, Aala, Thza, Thia, Bal, Fala, Pala, or X⁰-Phe;

AA⁶ is absent, the D- or L-isomer of R¹¹, an aromatic α -amino acid, F₅-Phe, Phe, Thr, Thr(Bzl), Ser, Ser(Bzl), or X⁰-Phe;


AA⁷ is absent, the D- or L-isomer of R¹¹ or the D- or L-isomer of an aromatic α -amino acid;

- 66 -

AA⁸ is a D- or L- isomer of R¹¹;

R¹ is H, E-, E(O)₂S-, E(O)C-, EOOC-, or R¹³;

R¹³ is a moiety of the formula



5 wherein R²¹ is (C₁₋₄)alkyl and s is 1, 2, 3, or 4;

X⁰ in the definition of AA² and AA⁵ is halogen, NO₂, OH, (C₁₋₆)alkyl, (C₁₋₆)alkoxy, mono- or di-(C₁₋₆)alkylamino, Bzl or O-Bzl;

X⁰ in the definition of AA⁶ is halogen, NO₂, OH, (C₁₋₆)alkyl, (C₁₋₆)alkoxy, mono- or di-(C₁₋₆)alkylamino, Bzl, O-Bzl, or NR⁹R¹⁰;

10 provided that:

at least one of AA¹ or AA² is present;

when AA¹ is absent, AA² and AA⁸ together form a bond; and


at least two of AA⁵, AA⁶, and AA⁷ are present;

or a pharmaceutically acceptable salt thereof.


15

5. A compound according to claim 2, wherein

AA¹ is absent, Ac-D-Phe, or the D- or L- isomer of R¹¹, Pip, Pro, or Ser, or of an aromatic α-amino acid selected from the group consisting of Cpa, Dip, Nal, Pal, and Phe;



20 AA² is absent, Aic, Pal, Phe, F₅-Phe, 4-NO₂-Phe, Trp, Tyr, Phe(4-O-Bzl)



- 67 -

- AA³ is the D- or L- isomer of an amino acid selected from the group consisting of Pen, Cys, hCys and Tmpa;
- AA⁴ is the D- or L-isomer of Trp, His, N-Me-Trp, β -Me-Trp, hTrp, or hHis;
- AA⁵ is Lys, hLys, N-Me-Lys, Orn, cis-4-Acha or 4-Pip-Ala;
- 5 AA⁶ is the D- or L-isomer of an amino acid selected from the group consisting of Cys, hCys, Pen and Tmpa;
- AA⁷ is A3c, A4c, A5c, A6c, Abu, Aic, β -Ala, Gaba, Nle, F₅-Phe, Phe, Pro, Sar, Ser, Thr, Thr(Bzl), Tyr, Val or absent; and
- AA⁸ is R¹¹, Nal, Thr, Thr(Bzl), Tyr, Phe(4-O-Bzl), or absent;
- 10 or a pharmaceutically acceptable salt thereof.

6. A compound according to claim 5, wherein
- AA¹ is absent or the D- or L- isomer of R¹¹, Pip or Pro, or of an aromatic α -amino acid selected from the group consisting of Cpa, Dip, Nal, Pal, Phe,
- 15 and Ac-Phe;
- AA² is Tyr, Pal, Phe, 4-NO₂-Phe, Trp, or absent;
- AA³ is a D- or L-isomer of Cys or Pen;
- AA⁴ is D-Trp;
- AA⁵ is Lys, Orn, or cis-4-Acha;
- 20 AA⁶ is a D- or L-isomer of Cys or Pen;
- AA⁷ is A3c, A4c, A5c, A6c, Abu, Aic, β -Ala, Gaba, Nle, Phe, Pro, Sar, Thr, Thr(Bzl), Tyr, Val, or absent; and
- AA⁸ is R¹¹, Thr, Tyr, Nal, or absent;
- or a pharmaceutically acceptable salt thereof.

25

7. A compound according to claim 3, wherein
- AA¹ is R¹¹, Aic, Hca, Pro, Ser, Ser(Bzl), Trp, Tyr, or a D- or L-isomer of an aromatic α -amino acid selected from the group consisting of Cpa, Nal, Ac-Nal, Phe, Ac-Phe, 4-NO₂-Phe, and Ac-4-NO₂-Phe;

- 68 -

- AA² is Pal, Phe, F₅-Phe, Tyr, or absent;
AA³ is a D- or L-isomer of Cys, hCys, Pen or Tmpa;
AA^{3b} is Pal, 4-Pal, His, Trp, Tyr, Phe(4-O-Bzl), Phe, or R¹¹;
AA⁴ is a D- or L-isomer of Trp or His;
- 5 AA⁵ is Lys, N-Me-Lys, Orn, hLys, cis-4-Acha, or 4-Pip-Ala;
AA⁶ is a D- or L-isomer of Cys, hCys, Pen or Tmpa;
AA⁷ is R¹¹, A4c, A5c, Abu, β -Ala, Gaba, Phe, F₅-Phe, Ser(Bzl), Thr, Thr(Bzl),
Phe(4-O-Bzl), or absent;
AA^{7b} is R¹¹, Nal, F₅-Phe, X⁰-Phe or absent, wherein X⁰ is halogen, NO₂, CH₃,
10 OH, Bzl or O-Bzl; and
AA⁸ is R¹¹, Nal, Tyr, Phe(4-O-Bzl), or absent;
or a pharmaceutically acceptable salt thereof.

8. A compound according to claim 7, wherein

- 15 AA¹ is R¹¹, Aic, Hca, Pro, Ser(Bzl), or a D- or L-isomer of an aromatic α -
amino acid selected from the group consisting of Cpa, Nal, Ac-Nal, Phe, Ac-
Phe, 4-NO₂-Phe, and Ac-4-NO₂-Phe;
AA² is Pal, Tyr, or absent;
AA³ is a D- or L-isomer of Cys or Pen;
- 20 AA^{3b} is R¹¹, Pal, 4-Pal, Trp, Tyr, Phe(4-O-Bzl), or Phe, wherein R¹¹ is (T)aeg;
AA⁴ is D-Trp;
AA⁵ is Lys, N-Me-Lys, Orn, or cis-4-Acha;
AA⁶ is a D- or L-isomer of Cys or Pen;
AA⁷ is R¹¹, A5c, Abu, Ser(Bzl), Thr, Thr(Bzl), Phe(4-O-Bzl), Gaba, or absent;
- 25 AA^{7b} is Nal, X⁰-Phe or absent; and
AA⁸ is Tyr or absent;
or a pharmaceutically acceptable salt thereof.

9. A compound according to claim 4, wherein

- 69 -

- AA¹ is Aic, Hyp, Cpa, D-Cpa, Nal, Pal, Phe, Pro, R¹¹, Tyr or absent;
AA² is Phe, Trp, F₅-Phe, His, Tyr, Phe(4-O-Bzl), or R¹¹;
AA³ is a D-isomer of Trp, His, or Pal;
AA⁴ is Lys, N-Me-Lys, Orn, hLys, cis-4-Acha, or 4-Pip-Ala;
5 AA⁵ is Pal, Phe(4-O-Bzl), Thr(Bzl), Thr, Sar, Gaba, β -Ala, A4c, A5c, A6c, Abu, Aic or absent;
AA⁶ is Thr, Tyr, Ser, F₅-Phe, Cpa, Nal, or D- or L-Phe;
AA⁷ is Nal, Pal, or absent; and
AA⁸ is R¹¹;
10 or a pharmaceutically acceptable salt thereof.

10. A compound according to claim 9, wherein
AA¹ is Cpa, Nal, Pal, Phe, Tyr or absent;
AA² is Phe, Tyr, Trp, or R¹¹;
15 AA³ is D-Trp;
AA⁴ is Lys, N-Me-Lys, or cis-4-Acha;
AA⁵ is Pal, Phe(4-O-Bzl), Aic, Gaba, A5c or absent;
AA⁶ is Thr, Nal, or D- or L-Phe;
AA⁷ is absent; and
20 AA⁸ is R¹¹;
or a pharmaceutically acceptable salt thereof.

11. A compound according to claim 2, wherein R¹ and R⁵ are absent
and the N-terminal amino acid and the C-terminal amino acid together form
25 an amide bond; or a pharmaceutically acceptable salt thereof.

12. A compound according to claim 3, wherein R¹ and R⁵ are absent
and the N-terminal amino acid and the C-terminal amino acid together form
an amide bond; or a pharmaceutically acceptable salt thereof.

13. A compound according to claim 6, wherein said compound is of the formula:

- Ac-D-Phe-Tyr-cyclo(D-Cys-D-Trp-Lys-Cys)-Abu-Thr-NH₂;
- 5 Nal-Tyr-cyclo(Cys-D-Trp-Lys-D-Cys)-Val-Nal-NH₂;
- Nal-Tyr-cyclo(Cys-D-Trp-Lys-D-Cys)-Abu-Nal-NH₂;
- D-Dip-Tyr-cyclo(Cys-D-Trp-Lys-D-Cys)-Abu-Nal-NH₂;
- Dip-Tyr-cyclo(D-Cys-D-Trp-Lys-D-Cys)-Abu-Nal-NH₂;
- Nal-Tyr-cyclo(D-Cys-D-Trp-Lys-D-Cys)-Abu-Nal-NH₂;
- 10 Dip-Tyr-cyclo(D-Cys-D-Trp-Lys-D-Cys)-Val-Nal-NH₂;
- Nal-Tyr-cyclo(D-Cys-D-Trp-Lys-D-Cys)-Val-Nal-NH₂;
- cyclo(D-Phe-Tyr-cyclo(D-Cys-D-Trp-Lys-Cys)-Abu-Thr);
- Cpa-Pal-cyclo(D-Cys-D-Trp-Lys-D-Cys)-A3c-Nal-NH₂;
- Cpa-Pal-cyclo(D-Cys-D-Trp-Lys-D-Cys)-A5c-Nal-NH₂;
- 15 Cpa-Pal-cyclo(D-Cys-D-Trp-Lys-D-Cys)-A6c-Nal-NH₂;
- (G(z))aeg-cyclo(D-Cys-D-Trp-Lys-D-Cys)-A5c-Nal-NH₂;
- Pal-cyclo(D-Cys-D-Trp-Lys-D-Cys)-A5c-Nal-NH₂;
- Cpa-Pal-cyclo(D-Cys-D-Trp-Lys-D-Cys)-β-Ala-Nal-NH₂;
- Cpa-Pal-cyclo(D-Cys-D-Trp-Lys-D-Cys)-Sar-Nal-NH₂;
- 20 Cpa-Pal-cyclo(D-Cys-D-Trp-Lys-D-Cys)-Gaba-Nal-NH₂;
- Cpa-Pal-cyclo(D-Cys-D-Trp-Lys-D-Cys)-Pro-Nal-NH₂;
- Pro-Phe-c(D-Cys-D-Trp-Lys-D-Cys)-Nle-Phe-NH₂;
- Pro-Phe-c(D-Cys-D-Trp-Lys-D-Cys)-Thr-Nle-NH₂;
- Pro-Phe-c(D-Cys-D-Trp-Lys-D-Cys)-Thr-Phe-NH₂;
- 25 Cpa-Phe-c(D-Cys-D-Trp-Lys-D-Cys)-Gaba-NH₂;
- Cpa-Phe-c(D-Cys-D-Trp-Lys-D-Cys)-Gaba-Tyr-NH₂;
- Pip-Phe-c(D-Cys-D-Trp-Lys-D-Cys)-NH₂;
- Pip-Phe-c(Cys-D-Trp-Lys-Cys)-Gaba-NH₂; or
- Pro-Phe-c(D-Cys-D-Trp-Lys-D-Cys)-Thr-NH₂;

or a pharmaceutically acceptable salt thereof.

14. A compound according to claim 6, wherein said compound is according to the formula:

- 5 Phe-cyclo(Cys-D-Trp-Lys-Cys)-Thr-NH₂;
Phe-Tyr-cyclo(D-Cys-D-Trp-Lys-Cys)-Abu-Thr-NH₂;
Ac-D-Phe-Tyr-cyclo(D-Cys-D-Trp-Lys-Cys)-Abu-Thr-NH₂;
Nal-Tyr-cyclo(Cys-D-Trp-Lys-D-Cys)-Val-Nal-NH₂;
Nal-Tyr-cyclo(Cys-D-Trp-Lys-D-Cys)-Abu-Nal-NH₂;
10 Dip-Tyr-cyclo(D-Cys-D-Trp-Lys-D-Cys)-Abu-Nal-NH₂;
Nal-Tyr-cyclo(D-Cys-D-Trp-Lys-D-Cys)-Abu-Nal-NH₂;
Dip-Tyr-cyclo(D-Cys-D-Trp-Lys-D-Cys)-Val-Nal-NH₂;
Nal-Tyr-cyclo(D-Cys-D-Trp-Lys-D-Cys)-Val-Nal-NH₂;
Cpa-Pal-cyclo(D-Cys-D-Trp-Lys-D-Cys)-A3c-Nal-NH₂;
15 Cpa-Pal-cyclo(D-Cys-D-Trp-Lys-D-Cys)-A5c-Nal-NH₂;
Cpa-Pal-cyclo(D-Cys-D-Trp-Lys-D-Cys)-A6c-Nal-NH₂;
(G(z))aeg-cyclo(D-Cys-D-Trp-Lys-D-Cys)-A5c-Nal-NH₂;
D-Cpa-cyclo(Cys-D-Trp-Lys-D-Cys)-A5c-Nal-NH₂;
Pal-cyclo(D-Cys-D-Trp-Lys-D-Cys)-A5c-Nal-NH₂;
20 Cpa-cyclo(D-Cys-D-Trp-Lys-D-Cys)-A5c-Nal-NH₂;
Cpa-Pal-cyclo(D-Cys-D-Trp-Lys-D-Cys)-β-Ala-Nal-NH₂;
Cpa-Pal-cyclo(D-Cys-D-Trp-Lys-D-Cys)-Sar-Nal-NH₂;
Cpa-Pal-cyclo(D-Cys-D-Trp-Lys-D-Cys)-Aic-Nal-NH₂;
Cpa-Pal-cyclo(D-Cys-D-Trp-Lys-D-Cys)-Gaba-Nal-NH₂;
25 Cpa-Pal-cyclo(D-Cys-D-Trp-Lys-D-Cys)-Pro-Nal-NH₂;
(T)aeg-cyclo(D-Cys-D-Trp-Lys-D-Cys)-(A)aeg-NH₂;
Cpa-Pal-cyclo(D-Cys-D-Trp-Lys-D-Cys)-A4c-Nal-NH₂;
Cpa-Pal-cyclo(D-Cys-D-Trp-Lys-D-Cys)-Nal-NH₂;
Pal-cyclo(D-Cys-D-Trp-Lys-D-Cys)-Nal-NH₂;

- 72 -

- Pro-Phe-cyclo(Cys-D-Trp-Lys-D-Cys)-Val-NH₂;
 Pro-Phe-cyclo(D-Cys-D-Trp-Lys-Cys)-Val-NH₂;
 Pip-4-NO₂-Phe-cyclo(D-Cys-D-Trp-Lys-D-Cys)-Nle-NH₂;
 (G)aeg-Pal-cyclo(D-Cys-D-Trp-Lys-D-Cys)-Thr(Bzl)-(C)aeg-NH₂; or
 5 (C)aeg-Pal-cyclo(D-Cys-D-Trp-Lys-D-Cys)-Thr(Bzl)-(G)aeg-NH₂;
 or a pharmaceutically acceptable salt thereof.

15. A compound according to claim 8, wherein said compound is according to the formula

- 10 Nal-cyclo(D-Cys-Tyr-D-Trp-Lys-Cys)-Nal-NH₂;
 D-Nal-cyclo(D-Cys-Tyr-D-Trp-Lys-Cys)-Nal-NH₂;
 D-Phe-cyclo(Cys-Tyr-D-Trp-Lys-Cys)-Thr-NH₂;
 D-4-NO₂-Phe-cyclo(D-Cys-Tyr-D-Trp-Lys-Cys)-Nal-NH₂;
 Ac-D-4-NO₂-Phe-cyclo(D-Cys-Tyr-D-Trp-Lys-Cys)-Nal-NH₂;
 15 D-4-NO₂-Phe-Pal-cyclo(D-Cys-Phe(4-O-Bzl)-D-Trp-Lys-Cys)-Tyr-NH₂;
 Cpa-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;
 D-4-NO₂-Phe-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Thr-Tyr-NH₂;
 D-4-NO₂-Phe-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Thr(Bzl)-NH₂;
 D-4-NO₂-Phe-cyclo(D-Cys-Pal-D-Trp-Lys-D-Cys)-Thr(Bzl)-Tyr-NH₂;
 20 D-4-NO₂-Phe-cyclo(D-Cys-Tyr-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;
 4-NO₂-Phe-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;
 D-Nal-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;
 Pro-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;
 Cpa-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Thr(Bzl)-Nal-NH₂;
 25 Ser(Bzl)-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Thr-Tyr-NH₂;
 (T)aeg-cyclo(D-Cys-Pal-D-Trp-Lys-D-Cys)-Thr(Bzl)-Tyr-NH₂;
 (A)aeg-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;
 (G)aeg-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;
 (T)aeg-cyclo(D-Cys-4-Pal-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;

- (T)aeg-cyclo(D-Cys-Tyr-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;
 (T)aeg-cyclo(D-Cys-Phe-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;
 (T)aeg-cyclo(D-Cys-(T)aeg-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;
 (T)aeg-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Ser(Bzl)-Tyr-NH₂;
 5 (T)aeg-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Phe(4-O-Bzl)-Tyr-NH₂;
 (T)aeg-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-A5c-Tyr-NH₂;
 (T)aeg-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Abu-Tyr-NH₂;
 D-Cpa-cyclo(D-Cys-(T)aeg-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;
 (C)aeg-c(D-Cys-Pal-D-Trp-Lys-D-Cys)-Thr(Bzl)-Tyr-NH₂;
 10 D-Cpa-c(D-Cys-Pal-D-Trp-Lys-D-Cys)Thr(Bzl)-Tyr-NH₂;
 (T)aeg-c(Pen-Pal-D-Trp-Lys-D-Cys)Thr(Bzl)-Tyr-NH₂;
 (T)aeg-c(D-Cys-Trp-D-Trp-Lys-D-Cys)Thr(Bzl)-Tyr-NH₂;
 (T)aeg-c(D-Cys-Phe-D-Trp-Lys-D-Cys)Thr(Bzl)-Tyr-NH₂;
 (T)aeg-c(D-Cys-Pal-D-Trp-Orn-D-Cys)Thr(Bzl)-Tyr-NH₂;
 15 (T)aeg-c(D-Cys-Pal-D-Trp-hLys-D-Cys)Thr(Bzl)-Tyr-NH₂;
 (T)aeg-c(D-Cys-Pal-D-Trp-Iamp-D-Cys)Thr(Bzl)-Tyr-NH₂;
 (T)aeg-c(D-Cys-Pal-D-Trp-Cha(4-am)-D-Cys)Thr(Bzl)-Tyr-NH₂;
 (T)aeg-c(D-Cys-Pal-D-Trp-Lys-D-Cys)-Ser(Bzl)-Tyr-NH₂;
 (T)aeg-c(D-Cys-Pal-D-Trp-Lys-D-Cys)Thr(Bzl)-D-Tyr-NH₂;
 20 (T)aeg-c(D-Cys-Pal-D-Trp-Lys-D-Cys)Thr(Bzl)-Trp-NH₂;
 (T)aeg-c(D-Cys-Pal-D-Trp-Lys-D-Pen)Thr(Bzl)-Tyr-NH₂;
 (C)aeg-c(D-Cys-Phe-D-Trp-Lys-D-Cys)Thr(Bzl)-Tyr-NH₂;
 Ina-c(D-Cys-Phe-D-Trp-Lys-D-Cys)-Thr(Bzl)-Tyr-NH₂;
 Mnf-c(D-Cys-Phe-D-Trp-Lys-D-Cys)-Thr(Bzl)-Tyr-NH₂;
 25 Inp-c(D-Cys-Phe-D-Trp-Lys-D-Cys)-Thr(Bzl)-Tyr-NH₂;
 Nua-c(D-Cys-Phe-D-Trp-Lys-D-Cys)-Thr(Bzl)-Tyr-NH₂;
 (T)aeg-Pal-c(D-Cys-D-Trp-Lys-D-Cys)Thr(Bzl)-Tyr-NH₂;
 (T)aeg-Pal-c(D-Cys-D-Trp-Lys-D-Cys)Tyr(Bzl)-Thr-NH₂;
 (C)aeg-Phe-c(D-Cys-D-Trp-Lys-D-Cys)Thr(Bzl)-Tyr-NH₂; or

- 74 -

(T)aeg-D-Trp-c(D-Cys-Pal-Lys-D-Cys)Thr(Bzl)-Leu-NH₂;
or a pharmaceutically acceptable salt thereof.

16. A compound according to claim 8, wherein said compound is
5 according to the formula

Hca-cyclo(D-Cys-Tyr-D-Trp-Lys-Cys)-Nal-NH₂;
Ac-Nal-cyclo(D-Cys-Tyr-D-Trp-Lys-Cys)-Nal-NH₂;
Ac-D-Phe-cyclo(D-Cys-Tyr-D-Trp-Lys-Cys)-Nal-NH₂;
Ac-D-Nal-cyclo(D-Cys-Tyr-D-Trp-Lys-Cys)-Nal-NH₂;
10 D-Phe-cyclo(D-Cys-Tyr-D-Trp-Lys-Cys)-Nal-NH₂;
Nal-cyclo(D-Cys-Tyr-D-Trp-Lys-Cys)-Nal-NH₂;
D-Nal-cyclo(D-Cys-Tyr-D-Trp-Lys-Cys)-Nal-NH₂;
D-Phe-cyclo(Cys-Tyr-D-Trp-Lys-Cys)-Thr-NH₂;
D-4-NO₂-Phe-cyclo(D-Cys-Tyr-D-Trp-Lys-Cys)-Nal-NH₂;
15 Ac-D-4-NO₂-Phe-cyclo(D-Cys-Tyr-D-Trp-Lys-Cys)-Nal-NH₂;
D-4-NO₂-Phe-Pal-cyclo(D-Cys-Phe(4-O-Bzl)-D-Trp-Lys-Cys)-Tyr-NH₂;
D-4-NO₂-Phe-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;
Cpa-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;
D-4-NO₂-Phe-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Thr(Bzl)-NH₂;
20 D-4-NO₂-Phe-cyclo(D-Cys-Pal-D-Trp-Lys-D-Cys)-Thr(Bzl)-Tyr-NH₂;
D-4-NO₂-Phe-cyclo(D-Cys-Tyr-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;
4-NO₂-Phe-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;
D-Nal-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;
Pro-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;
25 Cpa-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Thr(Bzl)-Nal-NH₂;
Ser(Bzl)-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Thr-Tyr-NH₂;
(T)aeg-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;
(C)aeg-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;
Aic-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;

- (C(z))aeg-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;
 (A(z))aeg-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;
 (T)aeg-cyclo(D-Cys-Pal-D-Trp-Lys-D-Cys)-Thr(Bzl)-Tyr-NH₂;
 (A)aeg-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;
 5 (G)aeg-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;
 (T)aeg-cyclo(D-Cys-4-Pal-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;
 (T)aeg-cyclo(D-Cys-Tyr-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;
 (T)aeg-cyclo(D-Cys-Phe-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;
 (T)aeg-cyclo(D-Cys-(T)aeg-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;
 10 (T)aeg-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Ser(Bzl)-Tyr-NH₂;
 (T)aeg-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Phe(4-O-Bzl)-Tyr-NH₂;
 (T)aeg-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-A5c-Tyr-NH₂;
 (T)aeg-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Abu-Tyr-NH₂;
 D-Cpa-cyclo(D-Cys-(T)aeg-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;
 15 (T)aeg-cyclo(D-Cys-Pal-D-Trp-Lys-D-Cys)-Thr(Bzl)-p-Me-Phe-NH₂;
 Ac-(T)aeg-cyclo(D-Cys-Pal-D-Trp-Lys-D-Cys)-Thr(Bzl)-Tyr-NH₂;
 (T)aeg-cyclo(D-Cys-Pal-D-Trp-Lys-D-Cys)-Nal-NH₂;
 D-Cpa-cyclo(D-Cys-Pal-D-Trp-Lys-D-Cys)-Nal-NH₂;
 (A)aeg-cyclo(D-Cys-Pal-D-Trp-Lys-D-Cys)-Thr(Bzl)-Tyr-NH₂; (C)aeg-
 20 cyclo(D-Cys-Pal-D-Trp-Lys-D-Cys)-Thr(Bzl)-Tyr-NH₂;
 (C)aeg-c(D-Cys-Pal-D-Trp-Lys-D-Cys)-Thr(Bzl)-Tyr-NH₂;
 D-Cpa-c(D-Cys-Pal-D-Trp-Lys-D-Cys)Thr(Bzl)-Tyr-NH₂;
 (T)aeg-c(Pen-Pal-D-Trp-Lys-D-Cys)Thr(Bzl)-Tyr-NH₂;
 (T)aeg-c(D-Cys-Trp-D-Trp-Lys-D-Cys)Thr(Bzl)-Tyr-NH₂;
 25 (T)aeg-c(D-Cys-Phe-D-Trp-Lys-D-Cys)Thr(Bzl)-Tyr-NH₂;
 (T)aeg-c(D-Cys-Pal-D-Trp-Orn-D-Cys)Thr(Bzl)-Tyr-NH₂;
 (T)aeg-c(D-Cys-Pal-D-Trp-hLys-D-Cys)Thr(Bzl)-Tyr-NH₂;
 (T)aeg-c(D-Cys-Pal-D-Trp-lamp-D-Cys)Thr(Bzl)-Tyr-NH₂;
 (T)aeg-c(D-Cys-Pal-D-Trp-Cha(4-am)-D-Cys)Thr(Bzl)-Tyr-NH₂;

- 76 -

- (T)aeg-c(D-Cys-Pal-D-Trp-Lys-D-Cys)-Ser(Bzl)-Tyr-NH₂;
 (T)aeg-c(D-Cys-Pal-D-Trp-Lys-D-Cys)Thr(Bzl)-D-Tyr-NH₂;
 (T)aeg-c(D-Cys-Pal-D-Trp-Lys-D-Cys)Thr(Bzl)-Trp-NH₂;
 (T)aeg-c(D-Cys-Pal-D-Trp-Lys-D-Pen)Thr(Bzl)-Tyr-NH₂;
 5 (C)aeg-c(D-Cys-Phe-D-Trp-Lys-D-Cys)Thr(Bzl)-Tyr-NH₂;
 Ina-c(D-Cys-Phe-D-Trp-Lys-D-Cys)-Thr(Bzl)-Tyr-NH₂;
 Mnf-c(D-Cys-Phe-D-Trp-Lys-D-Cys)-Thr(Bzl)-Tyr-NH₂;
 Inp-c(D-Cys-Phe-D-Trp-Lys-D-Cys)-Thr(Bzl)-Tyr-NH₂;
 Nua-c(D-Cys-Phe-D-Trp-Lys-D-Cys)-Thr(Bzl)-Tyr-NH₂;
 10 (T)aeg-Pal-c(D-Cys-D-Trp-Lys-D-Cys)Thr(Bzl)-Tyr-NH₂;
 (T)aeg-Pal-c(D-Cys-D-Trp-Lys-D-Cys)Tyr(Bzl)-Thr-NH₂;
 (C)aeg-Phe-c(D-Cys-D-Trp-Lys-D-Cys)Thr(Bzl)-Tyr-NH₂; or
 (T)aeg-D-Trp-c(D-Cys-Pal-Lys-D-Cys)Thr(Bzl)-Leu-NH₂;
 or a pharmaceutically acceptable salt thereof.

15

17. A compound according to claim 10, wherein said compound is according to the formula

- cyclo(Trp-D-Trp-Lys-Phe(4-O-Bzl)-Phe-(T)aeg);
 cyclo(Trp-D-Trp-Lys-Pal-Phe -(T)aeg); or
 20 cyclo(Phe-Phe-D-Trp-Lys-Thr-(T)aeg);
 or a pharmaceutically acceptable salt thereof.

18. A method of eliciting a neuromedin B receptor agonist effect in a subject in need thereof, wherein said method comprises administering to
 25 said subject an effective amount of a compound according to claim 13 or a pharmaceutically acceptable salt thereof.

19. A method of eliciting a somatostatin receptor agonist effect in a subject in need thereof, wherein said method comprises administering to

- 77 -

said subject an effective amount of a compound according to claim 14 or a pharmaceutically acceptable salt thereof.

20. A method of eliciting a neuromedin B receptor agonist effect in a
5 subject in need thereof, wherein said method comprises administering to said subject an effective amount of a compound according to claim 15 or a pharmaceutically acceptable salt thereof.

21. A method of eliciting a somatostatin receptor agonist effect in a
10 subject in need thereof, wherein said method comprises administering to said subject an effective amount of a compound according to claim 16 or a pharmaceutically acceptable salt thereof.

22. A method of eliciting a somatostatin receptor agonist effect in a
15 subject in need thereof, wherein said method comprises administering to said subject an effective amount of a compound according to claim 17 or a pharmaceutically acceptable salt thereof, provided said compound is not
cyclo(Trp-D-Trp-Lys-Phe(4-O-Bzl)-Phe-(T)aeg); or
cyclo(Trp-D-Trp-Lys-Pal-Phe -(T)aeg).

20

23. A method of eliciting a SSTR-1 agonist effect in a subject in need thereof, wherein said method comprises administering to said subject an effective amount of a compound according to claim 14 or a pharmaceutically acceptable salt thereof, provided said compound is not

25 Nal-Tyr-cyclo(Cys-D-Trp-Lys-D-Cys)-Val-Nal-NH₂;
Nal-Tyr-cyclo(Cys-D-Trp-Lys-D-Cys)-Abu-Nal-NH₂;
Dip-Tyr-cyclo(D-Cys-D-Trp-Lys-D-Cys)-Abu-Nal-NH₂;
Nal-Tyr-cyclo(D-Cys-D-Trp-Lys-D-Cys)-Abu-Nal-NH₂;
Dip-Tyr-cyclo(D-Cys-D-Trp-Lys-D-Cys)-Val-Nal-NH₂;

- 78 -

Nal-Tyr-cyclo(D-Cys-D-Trp-Lys-D-Cys)-Val-Nal-NH₂;
 Cpa-Pal-cyclo(D-Cys-D-Trp-Lys-D-Cys)-A3c-Nal-NH₂;
 Cpa-Pal-cyclo(D-Cys-D-Trp-Lys-D-Cys)-A5c-Nal-NH₂;
 Cpa-Pal-cyclo(D-Cys-D-Trp-Lys-D-Cys)-A6c-Nal-NH₂;
 5 (G(z))aeg-cyclo(D-Cys-D-Trp-Lys-D-Cys)-A5c-Nal-NH₂;
 D-Cpa-cyclo(Cys-D-Trp-Lys-D-Cys)-A5c-Nal-NH₂;
 Pal-cyclo(D-Cys-D-Trp-Lys-D-Cys)-A5c-Nal-NH₂;
 Cpa-cyclo(D-Cys-D-Trp-Lys-D-Cys)-A5c-Nal-NH₂;
 Cpa-Pal-cyclo(D-Cys-D-Trp-Lys-D-Cys)-β-Ala-Nal-NH₂;
 10 cyclo(D-Cys-D-Trp-Lys-D-Cys)-A5c-Nal-NH₂;
 Cpa-Pal-cyclo(D-Cys-D-Trp-Lys-D-Cys)-Sar-Nal-NH₂;
 Cpa-Pal-cyclo(D-Cys-D-Trp-Lys-D-Cys)-Aic-Nal-NH₂;
 Cpa-Pal-cyclo(D-Cys-D-Trp-Lys-D-Cys)-Gaba-Nal-NH₂; or
 Cpa-Pal-cyclo(D-Cys-D-Trp-Lys-D-Cys)-Pro-Nal-NH₂.

15

24. A method of eliciting a SSTR-1 agonist effect in a subject in need thereof, wherein said method comprises administering to said subject an effective amount of a compound according to claim 16 or a pharmaceutically acceptable salt thereof provided said compound is not

20 Ac-D-Phe-cyclo(D-Cys-Tyr-D-Trp-Lys-Cys)-Nal-NH₂;
 Ac-D-Nal-cyclo(D-Cys-Tyr-D-Trp-Lys-Cys)-Nal-NH₂;
 D-Phe-cyclo(D-Cys-Tyr-D-Trp-Lys-Cys)-Nal-NH₂;
 Nal-cyclo(D-Cys-Tyr-D-Trp-Lys-Cys)-Nal-NH₂;
 D-Nal-cyclo(D-Cys-Tyr-D-Trp-Lys-Cys)-Nal-NH₂;
 25 D-4-NO₂-Phe-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;
 Cpa-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;
 D-4-NO₂-Phe-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Thr(Bzl)-NH₂;
 D-4-NO₂-Phe-cyclo(D-Cys-Pal-D-Trp-Lys-D-Cys)-Thr(Bzl)-Tyr-NH₂;
 D-4-NO₂-Phe-cyclo(D-Cys-Tyr-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;

- 79 -

- 4-NO₂-Phe-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;
 D-Nal-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;
 Pro-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;
 Cpa-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Thr(Bzl)-Nal-NH₂;
 5 Ser(Bzl)-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Thr-Tyr-NH₂;
 (T)aeg-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;
 (C)aeg-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;
 Aic-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;
 (T)aeg-cyclo(D-Cys-Pal-D-Trp-Lys-D-Cys)-Thr(Bzl)-Tyr-NH₂;
 10 (A)aeg-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;
 (G)aeg-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;
 (T)aeg-cyclo(D-Cys-4-Pal-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;
 (T)aeg-cyclo(D-Cys-Tyr-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;
 (T)aeg-cyclo(D-Cys-Phe-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;
 15 (T)aeg-cyclo(D-Cys-(T)aeg-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂;
 (T)aeg-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Ser(Bzl)-Tyr-NH₂;
 (T)aeg-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Phe(4-O-Bzl)-Tyr-NH₂;
 (T)aeg-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-A5c-Tyr-NH₂;
 (T)aeg-cyclo(D-Cys-Pal-D-Trp-Lys-Cys)-Abu-Tyr-NH₂; or
 20 D-Cpa-cyclo(D-Cys-(T)aeg-D-Trp-Lys-Cys)-Thr(Bzl)-Tyr-NH₂.

25. A pharmaceutical composition comprising an effective amount of a compound according to claim 1 or a pharmaceutically acceptable salt thereof and a pharmaceutically acceptable carrier.

25

26. A method of treating a disease in a subject, said method comprising administering to said subject a therapeutically effective amount of a compound of claim 1, wherein said disease is selected from the list consisting of lung cancer, glioma, anorexia, hypothyroidism,

- 80 -

hyperaldosteronism, H. pylori proliferation, acromegaly, restenosis, Crohn's disease, systemic sclerosis, external and internal pancreatic pseudocysts and ascites, VIPoma, nesidoblastosis, hyperinsulinism, gastrinoma, Zollinger-Ellison Syndrome, diarrhea, AIDS related diarrhea, chemotherapy
5 related diarrhea, scleroderma, Irritable Bowel Syndrome, pancreatitis, small bowel obstruction, gastroesophageal reflux, duodenogastric reflux, Cushing's Syndrome, gonadotropinoma, hyperparathyroidism, Graves' Disease, diabetic neuropathy, Paget's disease, polycystic ovary disease, thyroid cancer, hepatome, leukemia, meningioma, cancer cachexia,
10 orthostatic hypotension, postprandial hypotension, panic attacks, GH secreting adenomas, Acromegaly, TSH secreting adenomas, prolactin secreting adenomas, insulinoma, glucagonoma, diabetes mellitus, hyperlipidemia, insulin insensitivity, Syndrome X, angiopathy, proliferative retinopathy, dawn phenomenon, Nephropathy, gastric acid secretion, peptic
15 ulcers, enterocutaneous fistula, pancreaticocutaneous fistula, Dumping syndrome, watery diarrhea syndrome, pancreatitis, gastrointestinal hormone secreting tumor, angiogenesis, arthritis, allograft rejection, graft vessel bleeding, portal hypertension, gastrointestinal bleeding, obesity, and opioid overdose.

20

COMBINED DECLARATION AND POWER OF ATTORNEY

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled NEUROMEDIN B AND SOMATOSTATIN RECEPTOR AGONISTS, the specification of which:

☐ is attached hereto.

☒ was filed on November 27, 2001 as Application Serial No. 09/980,133 and was amended on _____.

☒ was described and claimed in PCT International Application No. PCT/US00/15396 filed on June 5, 2000 and as amended under PCT Article 19 on _____.

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose all information I know to be material to patentability in accordance with Title 37, Code of Federal Regulations, §1.56.

I hereby claim the benefit under Title 35, United States Code, §119(e)(1) of any United States provisional application(s) listed below:

<u>U.S. Serial No.</u>	<u>Filing Date</u>	<u>Status</u>
60/137,655	6/04/1999	Expired

I hereby claim the benefit under Title 35, United States Code, §120 of any United States application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code, §112, I acknowledge the duty to disclose all information I know to be material to patentability as defined in Title 37, Code of Federal Regulations, §1.56(a) which became available between the filing date of the prior application and the national or PCT international filing date of this application:

<u>U.S. Serial No.</u>	<u>Filing Date</u>	<u>Status</u>
------------------------	--------------------	---------------

I hereby claim foreign priority benefits under Title 35, United States Code, §119 of any foreign application(s) for patent or inventor's certificate or of any PCT international application(s) designating at least one country other than the United States of America listed below and have also identified below any foreign application for patent or inventor's certificate or any PCT international application(s) designating at least one country other than the United States of America filed by me on the same subject matter having a filing date before that of the application(s) of which priority is claimed:

<u>Country</u>	<u>Application No.</u>	<u>Filing Date</u>	<u>Priority Claimed</u>
			<input type="checkbox"/> Yes <input type="checkbox"/> No
			<input type="checkbox"/> Yes <input type="checkbox"/> No

Combined Declaration and Power of Attorney

Page 2 of 2 Pages

I hereby appoint the following attorneys and/or agents to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith:

Y. Rocky Tsao, Reg. No. 34,053; Brian R. Morrill Reg. No. 42,098; Alan F. Feeney Reg. No. 43,609; Timothy A. French, Reg. No. 30,175; Eric L. Pahl, Reg. No. 32,590; Jeffrey D. Hsi, Reg. No. 40,024; and Frank R. Occhuiti, Reg. No. 35,306.

Address all telephone calls to Brian R. Morrill at telephone number (508) 478-0144.

Address all correspondence to Brian R. Morrill, Esq. at:

Biomeasure, Incorporated

27 Maple Street

Milford, MA 01757

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patents issued thereon.

Full Name of Inventor: BARRY A. MORGAN

Inventor's Signature: Barry A. Morgan

Date: 2.13.02

Residence Address: Franklin, Massachusetts

Citizenship: US

Post Office Address: 237 Prospect Street
Franklin, Massachusetts 02038
United States of America

Full Name of Inventor: DEAN SADAT-AALAE

Inventor's Signature: _____

Date: _____

Residence Address: _____

Citizenship: US

Post Office Address: 13940 Tahiti Way
Suite 221
Marina Del Ray, CA
90292
United States of America

Combined Declaration and Power of Attorney

Page 2 of 2 Pages

I hereby appoint the following attorneys and/or agents to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith:

Y. Rocky Tsao, Reg. No. 34,053; Brian R. Morrill Reg. No. 42,098; Alan F. Feeney Reg. No. 43,609; Timothy A. French, Reg. No. 30,175; Eric L. Pahl, Reg. No. 32,590; Jeffrey D. Hsi, Reg. No. 40,024; and Frank R. Occhuiti, Reg. No. 35,306.

Address all telephone calls to Brian R. Morrill at telephone number (508) 478-0144.

Address all correspondence to Brian R. Morrill, Esq. at:

Biomeasure, Incorporated
27 Maple Street
Milford, MA 01757

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patents issued thereon.

Full Name of Inventor: BARRY A. MORGAN

Inventor's Signature: _____ Date: _____
Residence Address: Franklin, Massachusetts
Citizenship: US
Post Office Address: 237 Prospect Street
Franklin, Massachusetts 02038
United States of America

Full Name of Inventor: DEAN SADAT-AALAE

Inventor's Signature: Dean Sadat-Aalae Date: 2/19/02
Residence Address: _____
Citizenship: US
Post Office Address: 13940 Tahiti Way
Suite 221
Marina Del Ray, CA
90292
United States of America